

Cellularity of free products of Boolean algebras (or topologies)

Saharon Shelah*

Institute of Mathematics
The Hebrew University of Jerusalem
91904 Jerusalem, Israel
and
Department of Mathematics
Rutgers University
New Brunswick, NJ 08854, USA

February 1, 2008

Abstract

The aim of this paper is to present an answer to Problem 1 of [M1], [M2]. We do this by proving in particular that

if μ is a strong limit singular cardinal, $\theta = (2^{\text{cf}(\mu)})^+$, $2^\mu = \mu^+$
then there are Boolean algebras $\mathbb{B}_1, \mathbb{B}_2$ such that $\text{c}(\mathbb{B}_1) = \mu$,
 $\text{c}(\mathbb{B}_2) < \theta$ but $\text{c}(\mathbb{B}_1 * \mathbb{B}_2) = \mu^+$.

Further we improve this result, deal with the method and the necessity of the assumptions.

*The research partially supported by “The Israel Science Foundation” administered by The Israel Academy of Sciences and Humanities. Publication 575. We thank Andrzej Roslanowski for writing sections 1 - 5 from lectures, 6 - 7 from notes

0 Introduction

Notation 0.1 1. In the present paper all cardinals are infinite so we will not repeat this additional demand. Cardinals will be denoted by λ, μ, θ (with possible indexes) while ordinal numbers will be called $\alpha, \beta, \zeta, \xi, \varepsilon, i, j$. Usually δ will stand for a limit ordinal (we may forget to repeat this assumption).

2. Sequences of ordinals will be called η, ν, ρ (with possible indexes). For sequences η_1, η_2 their longest common initial segment is denoted by $\eta_1 \wedge \eta_2$. The length of the sequence η is $\lg(\eta)$.

3. Ideals are supposed to be proper and contain all singletons. For a limit ordinal δ the ideal of bounded subsets of δ is denoted by J_δ^{bd} . If I is an ideal on a set X then I^+ is the family of I -large sets, i.e.

$$a \in I^+ \quad \text{if and only if} \quad a \subseteq X \text{ \& } a \notin I$$

and I^c is the dual filter of sets with the complements in I .

Notation 0.2 1. In a Boolean algebra we denote the Boolean operations by $\cap$ (and $\bigcap$), $\cup$ (and $\bigcup$), $-$. The distinguished elements are $\mathbf{0}$ and $\mathbf{1}$. In the cases which may be confusing we will add indexes to underline in which Boolean algebra the operation (the element) is considered, but generally we will not do it.

2. For a Boolean algebra $\mathbb{B}$ and an element $x \in \mathbb{B}$ we denote:

$$x^0 = x \quad \text{and} \quad x^1 = -x.$$

3. The free product of Boolean algebras $\mathbb{B}_1, \mathbb{B}_2$ is denoted by $\mathbb{B}_1 * \mathbb{B}_2$. We will use $\star$ to denote the free product of a family of Boolean algebras. For $\star_{i < \sigma} \mathbb{B}_i$ (wlog $\mathbb{B}_i$ pairwise disjoint) is the Boolean algebra generated by the formal intersection $\bar{b} = \bigcap_{i < \sigma} b_i$, $b_i \in \mathbb{B}$ freely except $\tau(\bar{b}^1, \dots, \bar{b}^n) = 0_{\mathbb{B}}$ iff some $u \in \sigma$ finite non empty for simplicity

$$\star_{i \in u} \mathbb{B}_i \models \tau\left(\bigcap_{i \in u} b_i^1 \dots\right) \neq 0$$

Definition 0.3 1. A Boolean algebra $\mathbb{B}$ satisfies the λ -cc if there is no family $\mathcal{F} \subseteq \mathbb{B}^+ \stackrel{\text{def}}{=} \mathbb{B} \setminus \{\mathbf{0}\}$ such that $|\mathcal{F}| = \lambda$ and any two members of $\mathcal{F}$ are disjoint (i.e. their meet in $\mathbb{B}$ is $\mathbf{0}$).

2. The cellularity of the algebra $\mathbb{B}$ is

$$\begin{aligned} c(\mathbb{B}) &= \sup\{|\mathcal{F}| : \mathcal{F} \subseteq \mathbb{B}^+ \ \& \ (\forall x, y \in \mathcal{F})(x \neq y \Rightarrow x \cap y = \mathbf{0})\}, \\ c^+(\mathbb{B}) &= \sup\{|\mathcal{F}|^+ : \mathcal{F} \subseteq \mathbb{B}^+ \ \& \ (\forall x, y \in \mathcal{F})(x \neq y \Rightarrow x \cap y = \mathbf{0})\}. \end{aligned}$$

3. For a topological space (X, τ) :

$$c(X, \tau) = \sup\{|\mathcal{U}| : \mathcal{U} \text{ is a family of pairwise disjoint nonempty open sets}\}.$$

The problem can be posed in each of the three ways (λ -cc is the way of forcing, the cellularity of Boolean algebras is the approach of Boolean algebraists, and the cellularity of a topological space is the way of general topologists). It is well known that the three are equivalent, though (1) makes the attainment problem more explicite. We use the second approach.

A stronger property than λ -cc is the λ -Knaster property. This property behaves nicely in free products – it is productive. We will use it in our construction.

Definition 0.4 A Boolean algebra $\mathbb{B}$ has the λ -Knaster property if for every sequence $\langle z_\varepsilon : \varepsilon < \lambda \rangle \subseteq \mathbb{B}^+$ there is $A \in [\lambda]^\lambda$ such that

$$\varepsilon_1, \varepsilon_2 \in A \quad \Rightarrow \quad z_{\varepsilon_1} \cap z_{\varepsilon_2} \neq \mathbf{0}.$$

We are interested in the behaviour of the cellularity of Boolean algebras when the free product of them is considered.

Thema 0.5 When, for Boolean algebras $\mathbb{B}_1, \mathbb{B}_2$

$$c^+(\mathbb{B}_1) \leq \lambda_1 \ \& \ c^+(\mathbb{B}_2) \leq \lambda_2 \quad \Rightarrow \quad c^+(\mathbb{B}_1 * \mathbb{B}_2) \leq \lambda_1 + \lambda_2?$$

There are a lot of results about it, particularly if $\lambda_1 = \lambda_2$ (see [Sh:g] or [M1], more [Sh 572]). It is well known that if

$$(\lambda_1^+ + \lambda_2^+) \longrightarrow (\lambda_1^+, \lambda_2^+)^2$$

then the answer is “yes”. These are exactly the cases for which “yes” answer is known. Under **GCH** the only problem which remained open was the one presented below:

The Problem We Address 0.6

(posed by D. Monk as Problem 1 in [M1], [M2] under GCH)

Are there Boolean algebras $\mathbb{B}_1, \mathbb{B}_2$ and cardinals μ, θ such that

1. $\lambda_1 = \mu$ is singular, $\mu > \lambda_2 = \theta > \text{cf}(\mu)$ and
2. $\text{c}(\mathbb{B}_1) = \mu$, $\text{c}(\mathbb{B}_2) \leq \theta$ but $\text{c}(\mathbb{B}_1 * \mathbb{B}_2) > \mu$?

We will answer this question proving in particular the following result (see 4.4):

If μ is a strong limit singular cardinal, $\theta = (2^{\text{cf}(\mu)})^+$, $2^\mu = \mu^+$ then there are Boolean algebras $\mathbb{B}_1, \mathbb{B}_2$ such that $\text{c}(\mathbb{B}_1) = \mu$, $\text{c}(\mathbb{B}_2) < \theta$ but $\text{c}(\mathbb{B}_1 * \mathbb{B}_2) = \mu^+$.

Later we deal with better results by refining the method.

Remark 0.7 On products of many Boolean algebras and square bracket arrows see [Sh 345], 1.2A, 1.3B.

If $\lambda \longrightarrow [\mu]^2$, $[\tau < \sigma \Rightarrow 2^\tau < \theta]$, the cardinals θ, σ are possibly finite, $\mathbb{B}_i$ (for $i < \theta$) are Boolean algebras such that for each $j < \theta$ the free product

$\star_{i \in \theta \setminus \{j\}} \mathbb{B}_i$ satisfies the μ -cc then the algebra $\mathbb{B} = \star_{i < \theta} \mathbb{B}_i$ satisfies the λ -cc.

[Why? if $\langle a_i^\zeta : i < \theta \rangle \in \prod_{i < \theta} \mathbb{B}_i^+$ for $\zeta < \lambda$ such that for every $\zeta < \xi < \lambda$ for some i , $\mathbb{B}_i \models "a_i^\zeta \cap a_i^\xi = \mathbf{0}"$, let $1 = 1(\zeta, \xi)$, and we can find $A \in [\lambda]^\mu$, $j < \theta$ such that for $\zeta < \xi$ from A , $1(\zeta, \xi) = j$, so $\langle a_i^\zeta : i < \theta, i \neq i^* \rangle$ for $\zeta \in A$ exemplifies $\star_{i \in \theta \setminus \{i^*\}} \mathbb{B}_i$ fails the μ -c.c.. We can deal also with ultraproducts and other products similarly.]

1 Preliminaries: products of ideals

Notation 1.1 For an ideal J on δ the quantifier $(\forall^J i < \delta)$ means “for all $i < \delta$ except a set from the ideal”, i.e.

$$(\forall^J i < \delta) \varphi(i) \equiv \{i < \delta : \neg \varphi(i)\} \in J.$$

The dual quantifier $(\exists^J i < \delta)$ means “for a J -positive set of $i < \delta$ ”.

Proposition 1.2 Assume that $\lambda^0 > \lambda^1 > \dots > \lambda^{n-1}$ are cardinals, I^ℓ are ideals on λ^ℓ (for $\ell < n$) and $B \subseteq \prod_{\ell < n} \lambda^\ell$. Further suppose that

$$(\alpha) \quad (\exists^{I^0} \gamma_0) \dots (\exists^{I^{n-1}} \gamma_{n-1}) (\langle \gamma_\ell : \ell < n \rangle \in B)$$

$$(\beta) \quad \text{the ideal } I^\ell \text{ is } (2^{\lambda^{\ell+1}})^+ \text{-complete (for } \ell + 1 < n \text{)}.$$

Then there are sets $X_\ell \subseteq \lambda^\ell$, $X_\ell \notin I^\ell$ such that $\prod_{\ell < n} X_\ell \subseteq B$.

[Note that this translates the situation to arity 1; it is a kind of polarized $(1, \dots, 1)$ -partitions with ideals.]

PROOF We show it by induction on n . Define

$$E_0 \stackrel{\text{def}}{=} \{(\gamma', \gamma'') : \gamma', \gamma'' < \lambda^0 \text{ and} \\ \text{for all } \gamma_1 < \lambda^1, \dots, \gamma_{n-1} < \lambda^{n-1} \text{ we have} \\ (\langle \gamma', \gamma_1, \dots, \gamma_{n-1} \rangle \in B \Leftrightarrow \langle \gamma'', \gamma_1, \dots, \gamma_{n-1} \rangle \in B)\}.$$

Clearly E_0 is an equivalence relation on λ^0 with $\leq 2^{\prod_{0 < m < n} \lambda^m} = 2^{\lambda^1}$ equivalence classes. Hence the set

$$A_0 \stackrel{\text{def}}{=} \bigcup \{A : A \text{ is an } E_0\text{-equivalence class, } A \in I^0\}$$

is in the ideal I^0 . Let

$$A_0^* \stackrel{\text{def}}{=} \{\gamma_0 < \lambda^0 : (\exists^{I^1} \gamma_1) \dots (\exists^{I^{n-1}} \gamma_{n-1}) (\langle \gamma_0, \gamma_1, \dots, \gamma_{n-1} \rangle \in B)\}.$$

The assumption (α) implies that $A_0^* \notin I^0$ and hence we may choose $\gamma_0^* \in A_0^* \setminus A_0$. Let

$$B_1 \stackrel{\text{def}}{=} \{\bar{\gamma} \in \prod_{k=1}^{n-1} \lambda^k : \langle \gamma_0^*, \bar{\gamma} \rangle \in B\}.$$

Since $\gamma_0^* \in A_0^*$ we are sure that

$$(\exists^{I^1} \gamma_1) \dots (\exists^{I^{n-1}} \gamma_{n-1}) (\langle \gamma_1, \dots, \gamma_{n-1} \rangle \in B_1).$$

Hence we may apply the inductive hypothesis for $n-1$ and B_1 and we find sets $X_1 \in (I^1)^+, \dots, X_{n-1} \in (I^{n-1})^+$ such that $\prod_{\ell=1}^{n-1} X_\ell \subseteq B_1$, so then

$$(\forall \gamma_1 \in X_1) \dots (\forall \gamma_{n-1} \in X_{n-1}) (\langle \gamma_0^*, \gamma_1, \dots, \gamma_{n-1} \rangle \in B).$$

Take X_0 to be the E_0 -equivalence class of γ_0^* (so $X_0 \in (I^0)^+$ as $\gamma_0^* \notin A_0$). By the definition of the relation E_0 and the choice of the sets X_ℓ we have that for each $\gamma_0 \in X_0$

$$(\forall \gamma_1 \in X_1) \dots (\forall \gamma_{n-1} \in X_{n-1}) (\langle \gamma_0, \gamma_1, \dots, \gamma_{n-1} \rangle \in B)$$

what means that $\prod_{\ell < n} X_\ell \subseteq B$. The proposition is proved. $\blacksquare$

Proposition 1.3 Assume that $\lambda_0 > \lambda_1 > \dots > \lambda_{n-1} \geq \sigma$ are cardinals, I_ℓ are ideals on λ_ℓ (for $\ell < n$) and $B \subseteq \prod_{\ell < n} \lambda_\ell$. Further suppose that

$$(\alpha) \quad (\exists^{I_0} \gamma_0) \dots (\exists^{I_{n-1}} \gamma_{n-1}) (\langle \gamma_\ell : \ell < n \rangle \in B)$$

$$(\beta) \quad \text{for each } \ell < n-1 \text{ the ideal } I_\ell \text{ is } ((\lambda_{\ell+1})^\sigma)^+ \text{-complete, } [\lambda_{n-1}]^{<\sigma} \subseteq I_{n-1}.$$

Then there are sets $X_\ell \in [\lambda_\ell]^\sigma$ such that $\prod_{\ell < n} X_\ell \subseteq B$.

PROOF The proof is by induction on n . If $n = 1$ then there is nothing to do as I_{n-1} contains all subsets of λ_{n-1} of size $< \sigma$ and $\lambda_{n-1} \geq \sigma$ so every $A \in I_{n-1}^+$ has cardinality $\geq \sigma$.

Let $n > 1$ and let

$$a_0 \stackrel{\text{def}}{=} \{\gamma \in \lambda_0 : (\exists^{I_1} \gamma_1) \dots (\exists^{I_{n-1}} \gamma_{n-1}) (\langle \gamma, \gamma_1, \dots, \gamma_{n-1} \rangle \in B)\}.$$

By our assumptions we know that $a_0 \in (I_0)^+$. For each $\gamma \in a_0$ we may apply the inductive hypothesis to the set

$$B_\gamma \stackrel{\text{def}}{=} \{\langle \gamma_1, \dots, \gamma_{n-1} \rangle \in \prod_{0 < \ell < n} \lambda_\ell : \langle \gamma, \gamma_1, \dots, \gamma_{n-1} \rangle \in B\}$$

and we get sets $X_1^\gamma \in [\lambda_1]^\sigma, \dots, X_{n-1}^\gamma \in [\lambda_{n-1}]^\sigma$ such that

$$\prod_{0 < \ell < n} X_\ell^\gamma \subseteq B_\gamma.$$

There is at most $(\lambda_1)^\sigma$ possible sequences $\langle X_1^\gamma, \dots, X_{n-1}^\gamma \rangle$, the ideal I_0 is $((\lambda_1)^\sigma)^+$ -complete so for some sequence $\langle X_1, \dots, X_{n-1} \rangle$ and a set $a^* \subseteq a_0$, $a^* \in (I_0)^+$ we have

$$(\forall \gamma \in a^*) (X_1^\gamma = X_1 \ \& \ \dots \ \& \ X_{n-1}^\gamma = X_{n-1}).$$

Choose $X_0 \in [a^*]^\sigma$ (remember that I_0 contains singletons and it is complete enough to make sure that $\sigma \leq |a^*|$). Clearly $\prod_{\ell < n} X_\ell \subseteq B$. ■

Remark 1.4 We can use $\sigma_0 \geq \sigma_1 \geq \dots \geq \sigma_{n-1}$, I_ℓ is $(\lambda_{\ell+1}^{\sigma_{\ell+1}})^+$ -complete, $[\lambda_\ell]^{<\sigma_\ell} \subseteq I_\ell$.

Proposition 1.5 Assume that $n < \omega$ and $\lambda_\ell^m, \chi_\ell^m, P_\ell^m, I_\ell^m, I^m$ and B are such that for $\ell, m \leq n$:

(α) I_ℓ^m is a χ_ℓ^m -complete ideal on λ_ℓ^m (for $\ell, m \leq n$)

(β) $P_\ell^m \subseteq \mathcal{P}(\lambda_\ell^m)$ is a family dense in $(I_\ell^m)^+$ in the sense that:

$$(\forall X \in (I_\ell^m)^+)(\exists a \in P_\ell^m)(a \subseteq X)$$

(γ) $I^m = \{X \subseteq \prod_{\ell \leq n} \lambda_\ell^m : \neg(\exists^{I_0^m} \gamma_0) \dots (\exists^{I_n^m} \gamma_n)(\langle \gamma_0, \dots, \gamma_n \rangle \in X)\}$

[thus I^m is the ideal on $\prod_{\ell \leq n} \lambda_\ell^m$ such that the dual filter $(I^m)^c$ is the Fubini product of filters $(I_0^m)^c, \dots, (I_n^m)^c$]

(δ) $\chi_{n-m}^m > \sum_{\ell=m+1}^n (|P_{n-\ell}^\ell| + \sum_{k=0}^{n-\ell} \lambda_k^\ell)$

(ε) $B \subseteq \prod_{m \leq n} \prod_{\ell \leq n} \lambda_\ell^m$ is a set satisfying

$$(\exists^{I^0} \eta_0)(\exists^{I^1} \eta_1) \dots (\exists^{I^n} \eta_n)(\langle \eta_0, \eta_1, \dots, \eta_n \rangle \in B).$$

Then there are sets $X_0, \dots, X_n$ such that for $m \leq n$:

(a) $X_m \subseteq \prod_{\ell \leq n-m} \lambda_\ell^m$

(b) if $\eta, \nu \in X_m$, $\eta \neq \nu$ then

(i) $\eta \upharpoonright (n-m) = \nu \upharpoonright (n-m)$

(ii) $\eta(n-m) \neq \nu(n-m)$

(c) $\{\eta(n-m) : \eta \in X_m\} \in P_{n-m}^m$ and

(d) for each $\langle \eta_0, \dots, \eta_n \rangle \in \prod_{m \leq n} X_m$ there is $\langle \eta_0^*, \dots, \eta_n^* \rangle \in B$ such that

$$(\forall m \leq n)(\eta_m \trianglelefteq \eta_m^*).$$

Remark 1.5.A:

1. Note that the sets X_m in the assertion of 1.5 may be thought of as sets of the form $X_m = \{\nu_m \hat{\ } \langle \alpha \rangle : \alpha \in a_m\}$ for some $\nu_m \in \prod_{\ell < n-m} \lambda_\ell^m$ and $a_m \in P_{n-m}^m$.
2. We will apply this proposition with $\lambda_\ell^m = \lambda_\ell$, $I_\ell^m = I_\ell$ and $\lambda_\ell > \chi_\ell > \sum_{k < \ell} \lambda_k$.

3. In the assumption (δ) of 1.5 we may have that the last sum on the right hand side of the inequality ranges from $k = 0$ to $n - \ell - 1$. We did not formulate that assumption in this way as with $n - \ell$ there it is easier to handle the induction step and this change is not important for our applications.
4. In the assertion **(d)** of 1.5 we can make η_ℓ^* depending on $\langle \eta_0, \dots, \eta_\ell \rangle$ only.

PROOF The proof is by induction on n . For $n = 0$ there is nothing to do. Let us describe the induction step.

Suppose $0 < n < \omega$ and $\lambda_\ell^m, \chi_\ell^m, P_\ell^m, I_\ell^m, I^m$ (for $\ell, m \leq n$) and B satisfy the assumptions (α) – (ε) . Let

$$B^* \stackrel{\text{def}}{=} \{ \langle \eta_0, \eta_1 \restriction n, \dots, \eta_n \restriction n \rangle : \eta_m \in \prod_{\ell \leq n} \lambda_\ell^m \text{ (for } m \leq n) \text{ and } \langle \eta_0, \eta_1, \dots, \eta_n \rangle \in B \}$$

and for $\eta_0 \in \prod_{\ell \leq n} \lambda_\ell^0$ let

$$B_{\eta_0}^* \stackrel{\text{def}}{=} \{ \langle \nu_1, \dots, \nu_n \rangle \in \prod_{m=1}^n \prod_{\ell=0}^{n-1} \lambda_\ell^m : \langle \eta_0, \nu_1, \dots, \nu_n \rangle \in B^* \}.$$

Let J^m (for $1 \leq m \leq n$) be the ideal on $\prod_{\ell=0}^{n-1} \lambda_\ell^m$ coming from the ideals I_ℓ^m , i.e. a set $X \subseteq \prod_{\ell < n} \lambda_\ell^m$ is in J^m if and only if

$$\neg(\exists^{I_0^m} \gamma_0) \dots (\exists^{I_{n-1}^m} \gamma_{n-1})(\langle \gamma_0, \dots, \gamma_{n-1} \rangle \in X).$$

Let us call the set $B_{\eta_0}^*$ *big* if

$$(\exists^{J^1} \nu_1) \dots (\exists^{J^n} \nu_n)(\langle \nu_1, \dots, \nu_n \rangle \in B_{\eta_0}^*).$$

We may write more explicitly what the bigness means: the above condition is equivalent to

$$\begin{aligned} & (\exists^{I_0^1} \gamma_0^1) \dots (\exists^{I_{n-1}^1} \gamma_{n-1}^1) \dots \\ & \dots (\exists^{I_0^n} \gamma_0^n) \dots (\exists^{I_{n-1}^n} \gamma_{n-1}^n)(\langle \langle \gamma_0^1, \dots, \gamma_{n-1}^1 \rangle, \dots, \langle \gamma_0^n, \dots, \gamma_{n-1}^n \rangle \rangle \in B_{\eta_0}^*) \end{aligned}$$

which means

$$\begin{aligned} & (\exists^{I_0^1} \gamma_0^1) \dots (\exists^{I_{n-1}^n} \gamma_{n-1}^n) \\ & (\exists \gamma_n^1) \dots (\exists \gamma_n^n)(\langle \eta_0, \langle \gamma_0^1, \dots, \gamma_n^1 \rangle, \dots, \langle \gamma_0^n, \dots, \gamma_n^n \rangle \rangle \in B). \end{aligned}$$

By the assumptions (γ) and (ε) we know that

$$\begin{aligned} & (\exists^{I_0^0} \gamma_0^0) \dots (\exists^{I_n^0} \gamma_n^0) (\exists^{I_0^1} \gamma_0^1) \dots (\exists^{I_n^1} \gamma_n^1) \dots \\ & \dots (\exists^{I_0^n} \gamma_0^n) \dots (\exists^{I_n^n} \gamma_n^n) (\langle \langle \gamma_0^0, \dots, \gamma_n^0 \rangle, \langle \gamma_0^1, \dots, \gamma_n^1 \rangle, \dots, \langle \gamma_0^n, \dots, \gamma_n^n \rangle \rangle \in B). \end{aligned}$$

Obviously any quantifier $(\exists^{I_\ell^m} \gamma_\ell^m)$ above may be replaced by $(\exists \gamma_\ell^m)$ and then “moved” right as much as we want. Consequently we get

$$\begin{aligned} & (\exists \gamma_0^0) \dots (\exists \gamma_{n-1}^0) (\exists^{I_n^0} \gamma_n^0) (\exists^{I_0^1} \gamma_0^1) \dots (\exists^{I_{n-1}^1} \gamma_{n-1}^1) \dots (\exists^{I_0^n} \gamma_0^n) \dots (\exists^{I_{n-1}^n} \gamma_{n-1}^n) \\ & (\exists \gamma_n^1) \dots (\exists \gamma_n^n) (\langle \langle \gamma_0^0, \dots, \gamma_n^0 \rangle, \langle \gamma_0^1, \dots, \gamma_n^1 \rangle, \dots, \langle \gamma_0^n, \dots, \gamma_n^n \rangle \rangle \in B) \end{aligned}$$

which means that

$$(\exists \gamma_0^0) \dots (\exists \gamma_{n-1}^0) (\exists^{I_n^0} \gamma_n^0) (B_{\langle \gamma_0^0, \dots, \gamma_n^0 \rangle}^* \text{ is big}).$$

Hence we find $\gamma_0^0, \dots, \gamma_{n-1}^0$ and a set $a \in (I_n^0)^+$ such that

$$(\forall \gamma \in a) (B_{\langle \gamma_0^0, \dots, \gamma_n^0 \rangle}^* \text{ is big}).$$

Note that the assumptions of the proposition are such that if we know that $B_{\eta_0}^*$ is big then we may apply the inductive hypothesis to

$$\lambda_\ell^m, \chi_\ell^m, P_\ell^m, I_\ell^m, J^m \quad (\text{for } 1 \leq m \leq n, \ell \leq n-1) \text{ and } B_{\eta_0}^*.$$

Consequently for each $\gamma \in a$ we find sets $X_1^\gamma, \dots, X_n^\gamma$ such that for $1 \leq m \leq n$:

$$(a)^* \quad X_m^\gamma \subseteq \prod_{\ell \leq n-m} \lambda_\ell^m$$

(b)* if $\eta, \nu \in X_m^\gamma$, $\eta \neq \nu$ then

$$(i) \quad \eta \upharpoonright (n-m) = \nu \upharpoonright (n-m)$$

$$(ii) \quad \eta(n-m) \neq \nu(n-m)$$

(c)* $\{\eta(n-m) : \eta \in X_m^\gamma\} \in P_{n-m}^m$ and

(d)* for all $\langle \eta_0, \dots, \eta_n \rangle \in \prod_{m \leq n} X_m^\gamma$ we have

$$(\exists \langle \eta_0^*, \dots, \eta_n^* \rangle \in B_{\langle \gamma_0^0, \dots, \gamma_{n-1}^0, \gamma \rangle}^*) (\forall 1 \leq m \leq n) (\nu_m \leq \nu_m^*).$$

Now we may ask how many possibilities for X_m^γ do we have: not too many. If we fix the common initial segment (see (b)*) the only freedom we have is

in choosing an element of P_{n-m}^m (see **(c)***). Consequently there are at most $|P_{n-m}^m| + \sum_{\ell \leq n-m} \lambda_\ell^m$ possible values for X_m^γ and hence there are at most

$$\sum_{m=1}^n (|P_{n-m}^m| + \sum_{\ell \leq n-m} \lambda_\ell^m) < \chi_n^0$$

possible values for the sequence $\langle X_1^\gamma, \dots, X_n^\gamma \rangle$. Since the ideal I_n^0 is χ_n^0 -complete we find a sequence $\langle X_1, \dots, X_n \rangle$ and a set $b \subseteq a$, $b \in (I_n^0)^+$ such that

$$(\forall \gamma \in b)(\langle X_1^\gamma, \dots, X_n^\gamma \rangle = \langle X_1, \dots, X_n \rangle).$$

Next choose $b_n^0 \in P_n^0$ such that $b_n^0 \subseteq b$ and put

$$X_0 = \{\langle \gamma_0^0, \dots, \gamma_{n-1}^0, \gamma \rangle : \gamma \in b_n^0\}.$$

Now it is a routine to check that the sets $X_0, X_1, \dots, X_n$ are as required (i.e. they satisfy clauses **(a)**–**(d)**). $\blacksquare$

2 Cofinal sequences in trees

Notation 2.1 1. For a tree $T \subseteq {}^{\delta>}\mu$ the set of δ -branches through T is

$$\lim_\delta(T) \stackrel{\text{def}}{=} \{\eta \in {}^\delta\mu : (\forall \alpha < \delta)(\eta \restriction \alpha \in T)\}.$$

The i -th level (for $i < \delta$) of the tree T is

$$T_i \stackrel{\text{def}}{=} T \cap {}^i\mu$$

$$\text{and } T_{<i} \stackrel{\text{def}}{=} \bigcup_{j < i} T_j.$$

If $\eta \in T$ then the set of immediate successors of η in T is

$$\text{succ}_T \stackrel{\text{def}}{=} \{\nu \in T : \eta \triangleleft \nu \text{ \& } \text{lg}(\nu) = \text{lg}(\eta) + 1\}.$$

We shall not distinguish strictly between $\text{succ}_T(\eta)$ and $\{\alpha : \eta \hat{\ } \langle \alpha \rangle \in T\}$.

Definition 2.2 1. $\mathcal{K}_{\mu, \delta}$ is the family of all pairs $(T, \bar{\lambda})$ such that $T \subseteq {}^{\delta>}\mu$ is a tree with δ levels and $\bar{\lambda} = \langle \lambda_\eta : \eta \in T \rangle$ is a sequence of cardinals such that for each $\eta \in T$ we have $\text{succ}_T(\eta) = \lambda_\eta$ (compare the previous remark about not distinguishing $\text{succ}_T(\eta)$ and $\{\alpha : \eta \hat{\ } \langle \alpha \rangle \in T\}$).

2. For a limit ordinal δ and a cardinal μ we let

$$\mathcal{K}_{\mu,\delta}^{\text{id}} \stackrel{\text{def}}{=} \{(T, \bar{\lambda}, \bar{I}) : (T, \bar{\lambda}) \in \mathcal{K}_{\mu,\delta}, \bar{I} = \langle I_\eta : \eta \in T \rangle \\ \text{each } I_\eta \text{ is an ideal on } \lambda_\eta = \text{succ}_T(\eta)\}.$$

Let $(T, \bar{\lambda}, \bar{I}) \in \mathcal{K}_{\mu,\delta}^{\text{id}}$ and let J be an ideal on δ (including J_δ^{bd} if we do not say otherwise). Further let $\bar{\eta} = \langle \eta_\alpha : \alpha < \lambda \rangle \subseteq \lim_\delta(T)$ be a sequence of δ -branches through T .

3. We say that $\bar{\eta}$ is J -cofinal in $(T, \bar{\lambda}, \bar{I})$ if

- (a) $\eta_\alpha \neq \eta_\beta$ for distinct $\alpha, \beta < \lambda$
- (b) for every sequence $\bar{A} = \langle A_\eta : \eta \in T \rangle \in \prod_{\eta \in T} I_\eta$ there is $\alpha^* < \lambda$ such that

$$\alpha^* \leq \alpha < \lambda \quad \Rightarrow \quad (\forall^J i < \delta)(\eta_\alpha \restriction (i+1) \notin A_{\eta_\alpha \restriction i}).$$

4. If I is an ideal on λ then we say that $(\bar{\eta}, I)$ is a J -cofinal pair for $(T, \bar{\lambda}, \bar{I})$ if

- (a) $\eta_\alpha \neq \eta_\beta$ for distinct $\alpha, \beta < \lambda$
- (b) for every sequence $\bar{A} = \langle A_\eta : \eta \in T \rangle \in \prod_{\eta \in T} I_\eta$ there is $A \in I$ such that

$$\alpha \in \lambda \setminus A \quad \Rightarrow \quad (\forall^J i < \delta)(\eta_\alpha \restriction (i+1) \notin A_{\eta_\alpha \restriction i}).$$

5. The sequence $\bar{\eta}$ is strongly J -cofinal in $(T, \bar{\lambda}, \bar{I})$ if

- (a) $\eta_\alpha \neq \eta_\beta$ for distinct $\alpha, \beta < \lambda$
- (b) for every $n < \omega$ and functions $F_0, \dots, F_n$ there is $\alpha^* < \lambda$ such that
if $m \leq n$, $\alpha_0 < \dots < \alpha_n < \lambda$, $\alpha^* \leq \alpha_m$
then the set

$$\{i < \delta : \begin{array}{l} (i) \ (\forall \ell < m)(\lambda_{\eta_{\alpha_\ell} \restriction i} < \lambda_{\eta_{\alpha_m} \restriction i}) \text{ and} \\ (ii) \ F_m(\eta_{\alpha_0} \restriction (i+1), \dots, \eta_{\alpha_{m-1}} \restriction (i+1), \eta_{\alpha_m} \restriction i, \dots, \eta_{\alpha_n} \restriction i) \in I_{\eta_{\alpha_m} \restriction i} \\ \text{(and well defined) but} \\ \eta_{\alpha_m} \restriction (i+1) \in F_m(\eta_{\alpha_0} \restriction (i+1), \dots, \eta_{\alpha_{m-1}} \restriction (i+1), \eta_{\alpha_m} \restriction i, \dots, \eta_{\alpha_n} \restriction i) \end{array}\}$$

is in the ideal J .

[Note: in (b) above we may have $\alpha^* < \alpha_0$, this causes no real change.]

6. The sequence $\bar{\eta}$ is stronger J -cofinal in $(T, \bar{\lambda}, \bar{I})$ if

- (a) $\eta_\alpha \neq \eta_\beta$ for distinct $\alpha, \beta < \lambda$
- (b) for every $n < \omega$ and functions $F_0, \dots, F_n$ there is $\alpha^* < \lambda$ such that
if $m \leq n$, $\alpha_0 < \dots < \alpha_n < \lambda$, $\alpha^* \leq \alpha_m$
then the set

$$\{i < \delta : \text{ (ii) } F_m(\eta_{\alpha_0} \upharpoonright (i+1), \dots, \eta_{\alpha_{m-1}} \upharpoonright (i+1), \eta_{\alpha_m} \upharpoonright i, \dots, \eta_{\alpha_n} \upharpoonright i) \in I_{\eta_{\alpha_m} \upharpoonright i} \\ \text{(and well defined) but} \\ \eta_{\alpha_m} \upharpoonright (i+1) \in F_m(\eta_{\alpha_0} \upharpoonright (i+1), \dots, \eta_{\alpha_{m-1}} \upharpoonright (i+1), \eta_{\alpha_m} \upharpoonright i, \dots, \eta_{\alpha_n} \upharpoonright i)\}$$

is in the ideal J .

7. The sequence $\bar{\eta}$ is strongest J -cofinal in $(T, \bar{\lambda}, \bar{I})$ if

- (a) $\eta_\alpha \neq \eta_\beta$ for distinct $\alpha, \beta < \lambda$
- (b) for every $n < \omega$ and functions $F_0, \dots, F_n$ there is $\alpha^* < \lambda$ such that
if $m \leq n$, $\alpha_0 < \dots < \alpha_n < \lambda$, $\alpha^* \leq \alpha_m$
then the set

$$\{i < \delta : \text{ (i')} \ (\exists \ell < m)(\lambda_{\eta_{\alpha_\ell} \upharpoonright i} \geq \lambda_{\eta_{\alpha_m} \upharpoonright i}) \text{ or} \\ \text{(ii) } F_m(\eta_{\alpha_0} \upharpoonright (i+1), \dots, \eta_{\alpha_{m-1}} \upharpoonright (i+1), \eta_{\alpha_m} \upharpoonright i, \dots, \eta_{\alpha_n} \upharpoonright i) \in I_{\eta_{\alpha_m} \upharpoonright i} \\ \text{(and well defined) but} \\ \eta_{\alpha_m} \upharpoonright (i+1) \in F_m(\eta_{\alpha_0} \upharpoonright (i+1), \dots, \eta_{\alpha_{m-1}} \upharpoonright (i+1), \eta_{\alpha_m} \upharpoonright i, \dots, \eta_{\alpha_n} \upharpoonright i)\}$$

is in the ideal J .

The sequence $\bar{\theta}$ is big* J -cofinal in $(T, \bar{\lambda}, \bar{I})$ if

- (a) $\eta_\alpha \neq \eta_\beta$ for distinct $\alpha, \beta < \lambda$
- (b) for every η and functions $F_0, \dots, F_n$ there is α^* such that
if $\alpha_0 < \dots < \alpha_n$ and $\alpha^* \leq \alpha_m$ then for $m \leq n$ the set

$$\{i < \delta : \text{ if } \nu_\ell = \begin{cases} \eta_{\alpha_\ell} \upharpoonright (i+1) & \text{if } \lambda_{\eta_{\alpha_\ell} \upharpoonright i} = \lambda_{\eta_{\alpha_{-m}} \upharpoonright i} \text{ or} \\ & \lambda_{\eta_{\alpha_\ell} \upharpoonright i} = \lambda_{\eta_{\alpha_n} \upharpoonright i} \text{ and } \eta_{\alpha_\ell}(i) < \eta_{\alpha_m}(i) \\ \eta_{\alpha_\ell} \upharpoonright i & \text{if not} \end{cases}$$

then we have

$$\eta_{\alpha_m}(i) \in F_m(\nu_\ell) \in I_{\eta_{\alpha_m} \upharpoonright i}\}$$

is in the ideal J .

We define “strongly* J -cofinal”, “stronger* J -cofinal” and “strongest* big J -cofinal” in almost the same way, replacing the requirement that $\alpha^* \leq \alpha_m$ in 5(b), 6(b), 7(b) above (respectively) by $\alpha^* \leq \alpha_0$.

- Remark 2.3** 1. Note that “strongest J -cofinal” implies “stronger J -cofinal” and this implies “strongly J -cofinal”. “Stronger J -cofinal” implies “ J -cofinal”. Also “bigger” $\Rightarrow$ “big” $\Rightarrow$ “cofinal”, “big” $\Rightarrow$ “strongly”.
2. The different notions of “strong J -cofinality” (the conditions (i) and (i’)) are to allow us to carry some diagonalization arguments.
3. The difference between “strongly J -cofinal” and “strongly* J -cofinal” etc is, in our context, immaterial. we may in all places in this paper replace the respective notion with its version with “*” and no harm will be done.

Remark 2.4 1. Remind **pcf**:

An important case is when $\langle \lambda_i : i < \delta \rangle$ is an increasing sequence of regular cardinals, $\lambda_i > \prod_{j < i} \lambda_j$, $\lambda_\eta = \lambda_{\text{lg}(\eta)}$, $I_\eta = J_{\lambda_\eta}^{\text{bd}}$ and $\lambda = \text{tcf}(\prod_{i < \delta} \lambda_i / J)$.

2. Moreover we are interested in more complicated I_η ’s (as in [Sh 430], section 5), connected to our problem, so “the existence of the true cofinality” is less clear. But the assumption $2^\mu = \mu^+$ will rescue us.
3. There are natural stronger demands of cofinality since here we are not interested just in x_α ’s but also in Boolean combinations. Thus naturally we are interested in behaviours of large sets of n -tuples, see 5.1.

Proposition 2.5 Suppose that $(T, \bar{\lambda}, \bar{I}) \in \mathcal{K}_{\mu, \delta}^{\text{id}}$, $\bar{\eta} = \langle \eta_\alpha : \alpha < \lambda \rangle \subseteq \lim_\delta(T)$ and J is an ideal on δ , $J \supseteq J_\delta^{\text{bd}}$.

1. Assume that

($\odot$) if $\alpha < \beta < \lambda$ then $(\forall^J i < \delta)(\lambda_{\eta_\alpha \restriction i} < \lambda_{\eta_\beta \restriction i})$.

Then the following are equivalent

“ $\bar{\eta}$ is strongly J -cofinal for $(T, \bar{\lambda}, \bar{I})$ ”

“ $\bar{\eta}$ is stronger J -cofinal for $(T, \bar{\lambda}, \bar{I})$ ”

“ $\bar{\eta}$ is strongest J -cofinal for $(T, \bar{\lambda}, \bar{I})$ ”.

“ $\bar{\eta}$ is big J -cofinal for $(T, \bar{\lambda}, \bar{I})$ ”.

2. If $I_\nu \supseteq J_{\lambda_\nu}^{\text{bd}}$ and $\lambda_\nu = \lambda_{\text{lg}(\nu)}$ for each $\nu \in T$ and the sequence $\bar{\eta}$ is stronger J -cofinal for $(T, \bar{\lambda}, \bar{I})$ then for some $\alpha^* < \lambda$ the sequence $\langle \eta_\alpha : \alpha^* \leq \alpha < \lambda \rangle$ is $<_J$ -increasing.
3. If $\eta \in T_i \Rightarrow \lambda_\eta = \lambda_i$ and $\bar{\eta} <_J$ -increasing in $\prod_{i < \delta}$ then “big” is equivalent to “stronger”. ■

Proposition 2.6 *Suppose that*

1. $\langle \lambda_i : i < \delta \rangle$ is an increasing sequence of regular cardinals, $\delta < \lambda_0$ is a limit ordinal,
2. $T = \bigcup_{i < \delta} \prod_{j < i} \lambda_j$, $I_\eta = I_{\text{lg}(\eta)} = J_{\lambda_{\text{lg}(\eta)}}^{\text{bd}}$, $\lambda_\eta = \lambda_{\text{lg}(\eta)}$,
3. J is an ideal on δ , $\lambda = \text{tcf}(\prod_{i < \delta} \lambda_i / J)$ and it is exemplified by a sequence $\bar{\eta} = \langle \eta_\alpha : \alpha < \lambda \rangle \subseteq \prod_{i < \delta} \lambda_i$
4. for each $i < \delta$

$$|\{\eta_\alpha \restriction i : \alpha < \lambda\}| < \lambda_i$$

(so e.g. $\lambda_i > \prod_{j < i} \lambda_j$ suffices).

Then the sequence $\bar{\eta}$ is J -cofinal in $(T, \bar{\lambda}, \bar{I})$.

PROOF First note that our assumptions imply that each ideal $I_\eta = I_{\text{lg}(\eta)}$ is $|\{\eta_\alpha \restriction \text{lg}(\eta) : \alpha < \lambda\}|^+$ -complete. Hence for each sequence $\bar{A} = \langle A_\eta : \eta \in T \rangle \in \prod_{\eta \in T} I_\eta$ and $i < \delta$ the set

$$A_i \stackrel{\text{def}}{=} \bigcup \{A_{\eta_\alpha \restriction i} : \alpha < \lambda\}$$

is in the ideal I_i , i.e. it is bounded in λ_i (for $i < \delta$). (We should remind here our convention which says in this case that we do not distinguish λ_i and $\text{succ}_T(\eta)$ if $\text{lg}(\eta) = i$, see 2.1.) Take $\eta^* \in \prod_{i < \delta} \lambda_i$ such that for each $i < \delta$ we have $A_i \subseteq \eta^*(i)$. As the sequence $\bar{\eta}$ realizes the true cofinality of $\prod_{i < \delta} \lambda_i / J$ we find $\alpha^* < \lambda$ such that

$$\alpha^* \leq \alpha < \lambda \quad \Rightarrow \quad \{i < \delta : \eta_\alpha(i) < \eta^*(i)\} \in J$$

which allows us to finish the proof. ■

It follows from the above proposition that the notion of J -cofinal sequences is not empty. Of course it is better to have “strongly (or even: stronger) J -cofinal” sequences $\bar{\eta}$. So it is nice to have that sometimes the weaker notion implies the stronger one.

Proposition 2.7 *Assume that δ is a limit ordinal, μ is a cardinal, $(T, \bar{\lambda}, \bar{I}) \in \mathcal{K}_{\mu, \delta}^{\text{id}}$. Let J be an ideal on δ such that $J \supseteq J_{\delta}^{\text{bd}}$ (which is our standard hypothesis). Further suppose that*

(*) *if $\eta \in T_i$ then the ideal I_{η} is $(|T_i| + \sum\{\lambda_{\nu} : \nu \in T_i \text{ \& } \lambda_{\nu} < \lambda_{\eta}\})^+$ -complete.*

Then each J -cofinal sequence $\bar{\eta}$ for $(T, \bar{\lambda}, \bar{I})$ is strongly J -cofinal for $(T, \bar{\lambda}, \bar{I})$.

If in addition $\eta \neq \nu \in T_i \Rightarrow \lambda_{\eta} \neq \lambda_{\nu}$ then $\bar{\eta}$ is big J -cofinal for $(T, \bar{\lambda}, \bar{I})$. Also if in addition

$\eta \in T_i \Rightarrow (\exists^{I_1} \nu \in T_i)(\lambda_{\nu} = \lambda_{\eta}) \vee [(\exists^{\leq \lambda_{\eta}} \nu \in T_i)(\lambda_{\nu} = \lambda_{\eta}) \text{ \& } I_{\eta} \text{ normal}]$

then $\bar{\eta}$ is big J -cofinal.

PROOF Let $n < \omega$ and $F_0, \dots, F_n$ be $(n+1)$ -place functions. First we define a sequence $\bar{A} = \langle A_{\eta} : \eta \in T \rangle$. For $m \leq n$ and a sequence $\langle \eta_m, \dots, \eta_n \rangle \subseteq T_i$ we put

$$A_{\langle \eta_m, \dots, \eta_n \rangle}^m = \bigcup \{ F_m(\nu_0, \dots, \nu_{m-1}, \eta_m, \dots, \eta_n) : \nu_0, \dots, \nu_{m-1} \in T_{i+1}, \\ (\nu_0, \dots, \nu_{m-1}, \eta_m, \dots, \eta_n) \in \text{dom}(F), \\ \lambda_{\nu_0 \upharpoonright i} < \lambda_{\eta}, \dots, \lambda_{\nu_{m-1} \upharpoonright i} < \lambda_{\eta_m} \\ \text{and } F(\nu_0, \dots, \nu_{m-1}, \eta_m, \dots, \eta_n) \in I_{\eta_m} \}$$

and next for $\eta \in T_i$ let

$$A_{\eta} = \bigcup \{ A_{\langle \eta, \eta_{m+1}, \dots, \eta_n \rangle}^m : m \leq n \text{ \& } \eta_{m+1}, \dots, \eta_n \in T_i \}.$$

Note that the assumption (*) was set up so that $A_{\langle \eta_m, \dots, \eta_n \rangle}^m \in I_{\eta_m}$ and the sets A_{η} are in I_{η} (for $\eta \in T$).

By the J -cofinality of $\bar{\eta}$, for some $\alpha^* < \lambda$ we have

$$\alpha^* \leq \alpha < \lambda \Rightarrow (\forall^J i < \delta)(\eta_{\alpha} \upharpoonright (i+1) \notin A_{\eta_{\alpha} \upharpoonright i}).$$

We are going to prove that this α^* is as required in the definition of strongly J -cofinal sequences. So suppose that $m \leq n$, $\alpha_0 < \dots < \alpha_n < \lambda$ and $\alpha^* \leq \alpha_m$. By the choice of α^* we have that the set $A \stackrel{\text{def}}{=} \{i < \delta : \eta_{\alpha_m} \upharpoonright (i+1) \in A_{\eta_{\alpha_m} \upharpoonright i}\}$ is in the ideal J . But if $i < \delta$ is such that

$$(\forall \ell < m)(\lambda_{\eta_{\alpha_{\ell}}} \upharpoonright i < \lambda_{\eta_{\alpha_m} \upharpoonright i}) \text{ \& } \\ F(\eta_{\alpha_0} \upharpoonright (i+1), \dots, \eta_{\alpha_{m-1}} \upharpoonright (i+1), \eta_{\alpha_m} \upharpoonright i, \dots, \eta_{\alpha_n} \upharpoonright i) \in I_{\eta_{\alpha_m} \upharpoonright i} \text{ but } \\ \eta_{\alpha_m} \upharpoonright (i+1) \in F(\eta_{\alpha_0} \upharpoonright (i+1), \dots, \eta_{\alpha_{m-1}} \upharpoonright (i+1), \eta_{\alpha_m} \upharpoonright i, \dots, \eta_{\alpha_n} \upharpoonright i)$$

then clearly $\eta_{\alpha_m} \restriction (i+1) \in A_{\langle \eta_{\alpha_m} \restriction i, \dots, \eta_{\alpha_n} \restriction i \rangle}^m$ and so $i \in A$. This finishes the proof.

The “big” version should be clear too. ■

Proposition 2.8 *Assume that μ is a strong limit uncountable cardinal and $\langle \mu_i : i < \delta \rangle$ is an increasing sequence of cardinals with limit μ . Further suppose that $(T, \bar{\lambda}, \bar{I}) \in \mathcal{K}_{\mu, \delta}^{\text{id}}$, $|T_i| \leq \mu_i$ (for $i < \delta$), $\lambda_\eta < \mu$ and each I_η is $\mu_{\text{lg}(\eta)}^+$ -complete and contains all singletons (for $\eta \in T$). Finally assume $2^\mu = \mu^+$ and let J be an ideal on δ , $J \supseteq J_\delta^{\text{bd}}$.*

Then there exists a stronger J -cofinal sequence $\bar{\eta}$ for $(T, \bar{\lambda}, \bar{I})$ of the length μ^+ (even for $J = J_\delta^{\text{bd}}$).

We can get “big” if

$$\rho \neq \eta \in T_i \ \& \ \lambda_\rho = \lambda_\eta \Rightarrow (\exists^{\leq \lambda_\eta} \nu \in T_i)(\lambda_\nu = \lambda_\eta) \ \& \ I_\eta \text{ normal}.$$

PROOF This is a straight diagonal argument. Put

$$Y \stackrel{\text{def}}{=} \{ \langle F_0, \dots, F_n \rangle : n < \omega \text{ and each } F_\ell \text{ is a function with} \\ \text{dom}(F) \subseteq T^{n+1}, \text{rng}(F) \subseteq \bigcup_{\eta \in T} I_\eta \}.$$

Since $|Y| = \mu^\mu = \mu^+$ (remember that μ is strong limit and $\lambda_\eta < \mu$ for $\eta \in T$) we may choose an enumeration $Y = \{ \langle F_0^\xi, \dots, F_{n_\xi}^\xi \rangle : \xi < \mu^+ \}$. For each $\zeta < \mu^+$ choose an increasing sequence $\langle \mathcal{A}_i^\zeta : i < \delta \rangle$ such that $|\mathcal{A}_i^\zeta| \leq \mu_i$ and $\zeta = \bigcup_{i < \delta} \mathcal{A}_i^\zeta$. Now we choose by induction on $\zeta < \mu^+$ branches η_ζ such that for each ζ the restriction $\eta_\zeta \restriction i$ is defined by induction on ζ as follows. If $i = 0$ or i is limit then there is nothing to do. Suppose now that we have defined $\eta_\zeta \restriction i$ and η_ξ for $\xi < \zeta$. We find $\eta_\zeta(i)$ such that

$$(\alpha) \ \eta_\zeta(i) \in \lambda_{\eta_\zeta \restriction i}$$

$$(\beta) \text{ if } \varepsilon \in \mathcal{A}_i^\zeta, m \leq n_\varepsilon, \alpha_0, \dots, \alpha_{m-1} \in \mathcal{A}_i^\zeta \text{ (hence } \alpha_\ell < \zeta \text{ so } \eta_{\alpha_\ell} \text{ are defined already), } \nu_{m+1}, \dots, \nu_n \in T_i \text{ and}$$

$$F_m^\varepsilon(\eta_{\alpha_0} \restriction (i+1), \dots, \eta_{\alpha_{m-1}} \restriction (i+1), \eta_\zeta \restriction i, \nu_{m+1}, \dots, \nu_n) \in I_{\eta_\zeta \restriction i} \text{ and well defined}$$

then

$$\eta_\zeta \restriction (i+1) \notin F_m^\varepsilon(\eta_{\alpha_0} \restriction (i+1), \dots, \eta_{\alpha_{m-1}} \restriction (i+1), \eta_\zeta \restriction i, \nu_{m+1}, \dots, \nu_n)$$

$$(\gamma) \ \eta_\zeta \restriction (i+1) \notin \{ \eta_\varepsilon \restriction (i+1) : \varepsilon \in \mathcal{A}_i^\zeta \}.$$

Why it is possible? Note that there is $\leq \aleph_0 + |\mathcal{A}_i^\zeta| + |\mathcal{A}_i^\zeta|^{<\aleph_0} + |T_i| \leq \mu_i$ negative demands and each of them says that $\eta_\zeta \restriction (i+1)$ is not in some set from $I_{\eta_\zeta \restriction i}$ (remember that we have assumed that the ideals $I_{\eta_\zeta \restriction i}$ contain singletons). Consequently using the completeness of the ideal we may satisfy the requirements (α) – (γ) above.

Now of course $\eta_\zeta \in \lim_\delta(T)$. Moreover if $\varepsilon < \zeta < \mu^+$ then $(\exists i < \delta)(\varepsilon \in \mathcal{A}_i^\zeta)$ which implies $(\exists i < \delta)(\eta_\varepsilon \restriction (i+1) \neq \eta_\zeta \restriction (i+1))$. Consequently

$$\varepsilon < \zeta < \mu^+ \Rightarrow \eta_\varepsilon \neq \eta_\zeta.$$

Checking the demand **(b)** of “stronger J -cofinal” is straightforward: for functions $F_0, \dots, F_n$ (and $n \in \omega$) take ε such that

$$\langle F_0, \dots, F_n \rangle = \langle F_0^\varepsilon, \dots, F_n^\varepsilon \rangle$$

and put $\alpha^* = \varepsilon + 1$. Suppose now that $m \leq n$, $\alpha_0 < \dots < \alpha_n < \lambda$, $\alpha^* \leq \alpha_m$. Let $i^* < \delta$ be such that for $i > i^*$ we have

$$\varepsilon, \alpha_0, \dots, \alpha_{m-1} \in \mathcal{A}_i^{\alpha_m}.$$

Then by the choice of $\eta_{\alpha_m} \restriction (i+1)$ we have that for each $i > i^*$

if $F_m^\varepsilon(\eta_{\alpha_0} \restriction (i+1), \dots, \eta_{\alpha_{m-1}} \restriction (i+1), \eta_\zeta \restriction i, \eta_{\alpha_{m+1}} \restriction i, \dots, \eta_{\alpha_n} \restriction i) \in I_{\eta_{\alpha_m} \restriction i}$
then $\eta_{\alpha_m} \restriction i \notin F_m^\varepsilon(\eta_{\alpha_0} \restriction (i+1), \dots, \eta_{\alpha_{m-1}} \restriction (i+1), \eta_\zeta \restriction i, \eta_{\alpha_{m+1}} \restriction i, \dots, \eta_{\alpha_n} \restriction i)$.

This finishes the proof. ■

Remark 2.9 The proof above can be carried for functions F which depend on $(\eta_{\alpha_0}, \dots, \eta_{\alpha_{m-1}}, \eta_{\alpha_m} \restriction i, \dots, \eta_{\alpha_n} \restriction i)$. This will be natural later.

Let us note that if the ideals I_η are sufficiently complete then J -cofinal sequences cannot be too short.

Proposition 2.10 *Suppose that $(T, \bar{\lambda}, \bar{I}) \in \mathcal{K}_{\mu, \delta}^{\text{id}}$ is such that for each $\eta \in T_i$, $i < \delta$ the ideal I_η is $(\kappa_i)^+$ -complete (enough if $[\lambda_\eta]^{\kappa_i} \subseteq I_\eta$). Let $J \supseteq J_\delta^{\text{bd}}$ be an ideal on δ and let $\bar{\eta} = \langle \eta_\alpha : \alpha < \delta^* \rangle$ be a J -cofinal sequence for $(T, \bar{\lambda}, \bar{I})$. Then*

$$\delta^* > \limsup_J \kappa_i$$

and consequently

$$\text{cf}(\delta^*) > \limsup_J \kappa_i.$$

PROOF Fix an enumeration $\delta^* = \{\alpha_\varepsilon : \varepsilon < |\delta^*|\}$ and for $\alpha < \delta^*$ let $\zeta(\alpha)$ be the unique ζ such that $\alpha = \alpha_\zeta$.
For $\eta \in T_i$, $i < \delta$ put

$$A_\eta \stackrel{\text{def}}{=} \{\nu \in \text{succ}_T(\eta) : (\exists \varepsilon \leq \kappa_i)(\nu \triangleleft \eta_\varepsilon)\}.$$

Clearly $|A_\eta| \leq \kappa_i$ and hence $A_\eta \in I_\eta$. Apply the J -cofinality of $\bar{\eta}$ to the sequence $A = \langle A_\eta : \eta \in T \rangle$. Thus there is $\alpha^* < \delta^*$ such that for each $\alpha \in [\alpha^*, \delta^*)$ we have

$$(\forall^J i < \delta)(\eta_\alpha \restriction (i+1) \notin A_{\eta_\alpha \restriction i})$$

and hence

$$(\forall^J i < \delta)(\zeta(\alpha) > \kappa_i)$$

and consequently

$$\zeta(\alpha) \geq \limsup_J \kappa_i.$$

Hence we conclude that $|\delta^*| > \limsup_J \kappa_i$.

For the part “consequently” of the proposition note that if $\langle \eta_\alpha : \alpha < \delta^* \rangle$ is J -cofinal (in $(T, \bar{\lambda}, \bar{I})$) and $A \subseteq \delta^*$ is cofinal in δ^* then $\langle \eta_\alpha : \alpha \in A \rangle$ is J -cofinal too. ■

Remark 2.11 1. So if we have a J -cofinal sequence of the length δ^* then we also have one of the length $\text{cf}(\delta^*)$. Thus assuming regularity of the length is natural.

2. Moreover the assumption that the length of the sequence is above $|\delta| + |T|$ is very natural and in most cases it will follow from the J -cofinality (and completeness assumptions). However we will try to state this condition in the assumptions whenever it is used in the proof (even if it can be concluded from the other assumptions).

3 Getting $(\kappa, \text{not } \lambda)$ -Knaster algebras

Proposition 3.1 Let λ, σ be cardinals such that $(\forall \alpha < \sigma)(2^{|\alpha|} < \lambda)$, σ is regular. Then there are a Boolean algebra $\mathbb{B}$, a sequence $\langle y_\alpha : \alpha < \lambda \rangle \subseteq \mathbb{B}^+$ and an ideal I on λ such that

(a) if $X \subseteq \lambda$, $X \notin I$ then $(\exists \alpha, \beta \in X)(\mathbb{B} \models y_\alpha \cap y_\beta = \mathbf{0})$

(b) the ideal I is σ -complete

(c) the algebra $\mathbb{B}$ satisfies the μ -Knaster condition for any regular uncountable μ (really $\mathbb{B}$ is free).

PROOF Let $\mathbb{B}$ be the Boolean algebra freely generated by $\{z_\alpha : \alpha < \lambda\}$ (so the demand (c) is satisfied). Let $A = \{(\alpha, \beta) : \alpha < \beta < \lambda\}$ and $y_{(\alpha, \beta)} = z_\alpha - z_\beta (\neq \mathbf{0})$ (for $(\alpha, \beta) \in A$). The ideal I of subsets of A defined by

a set $X \subseteq A$ is in I if and only if
 there are $\zeta < \sigma$, $X_\varepsilon \subseteq A$ (for $\varepsilon < \zeta$) such that $X \subseteq \bigcup_{\varepsilon < \zeta} X_\varepsilon$ and
 for every $\varepsilon < \zeta$ no two $y_{(\alpha_1, \beta_1)}, y_{(\alpha_2, \beta_2)} \in X_\varepsilon$ are disjoint in $\mathbb{B}$.

First note that

Claim 3.1.1 $A \notin I$.

Why? If not then we have witnesses $\zeta < \sigma$ and X_ε (for $\varepsilon < \zeta$) for it. So $A = \bigcup_{\varepsilon < \zeta} X_\varepsilon$ and hence for $(\alpha, \beta) \in A$ we have $\varepsilon(\alpha, \beta)$ such that $y_{(\alpha, \beta)} \in X_{\varepsilon(\alpha, \beta)}$.

So $\varepsilon(\cdot, \cdot)$ is actually a function from $[\lambda]^2$ to $\zeta < \sigma$. By the Erdős–Rado theorem we find $\alpha < \beta < \gamma < \lambda$ such that $\varepsilon(\alpha, \beta) = \varepsilon(\beta, \gamma)$. But

$$y_{(\alpha, \beta)} \cap y_{(\beta, \gamma)} = (z_\alpha - z_\beta) \cap (z_\beta - z_\gamma) = \mathbf{0}$$

so $(\alpha, \beta), (\beta, \gamma)$ cannot be in the same X_ε – a contradiction.

To finish the proof note that I is σ -complete (as σ is regular), if $X \notin I$ then, by the definition of I , there are two disjoint elements in $\{y_{(\alpha, \beta)} : (\alpha, \beta) \in X\}$. Finally $|A| = \lambda$. ■

Definition 3.2 (a) A pair $(\mathbb{B}, \bar{y})$ is called a λ -marked Boolean algebra if $\mathbb{B}$ is a Boolean algebra and $\bar{y} = \langle y_\alpha : \alpha < \lambda \rangle$ is a sequence of non-zero elements of $\mathbb{B}$.

(b) A triple $(\mathbb{B}, \bar{y}, I)$ is called a (λ, χ) -well marked Boolean algebra if $(\mathbb{B}, \bar{y})$ is a λ -marked Boolean algebra, χ is a regular cardinal and I is a (proper) χ -complete ideal on λ such that

$$\{A \subseteq \lambda : (\forall \alpha, \beta \in A)(\mathbb{B} \models y_\alpha \cap y_\beta \neq \mathbf{0})\} \subseteq I.$$

By λ -well marked Boolean algebra we will mean $(\lambda, \aleph_0)$ -well marked one. As in the above situation λ can be read from $\bar{y}$ (as $\lambda = \text{lg}(\bar{y})$) we may omit it and then we may speak just about well marked Boolean algebras.

Remark 3.3 Thus proposition 3.1 says that if λ, σ are regular cardinals and

$$(\forall \alpha < \sigma)(2^{|\alpha|} < \lambda)$$

then there exists a (λ, σ) -well marked Boolean algebra $(\mathbb{B}, \bar{y}, I)$ such that $\mathbb{B}$ satisfies the κ -Knaster property for every κ .

Definition 3.4 (a) For cardinals μ and λ and a limit ordinal δ , a (δ, μ, λ) -constructor is a system

$$\mathcal{C} = (T, \bar{\lambda}, \bar{\eta}, \langle (\mathbb{B}_\eta, \bar{y}_\eta) : \eta \in T \rangle)$$

such that

1. $(T, \bar{\lambda}) \in \mathcal{K}_{\mu, \delta}$,
2. $\bar{\eta} = \langle \eta_i : i \in \lambda \rangle$ where $\eta_i \in \lim_\delta(T)$ (for $i < \lambda$) are distinct δ -branches through T and
3. for each $\eta \in T$: $(\mathbb{B}_\eta, \bar{y}_\eta)$ is a λ_η -marked Boolean algebra, i.e. $\bar{y}_\eta = \langle y_{\eta \restriction \langle \alpha \rangle} : \alpha < \lambda_\eta \rangle \subseteq \mathbb{B}_\eta^+$ (usually this will be an enumeration of $\mathbb{B}_\eta^+$).

(b) Let $\mathcal{C}$ be a constructor (as above). We define Boolean algebras $\mathbb{B}_2 = \mathbb{B}^{\text{red}} = \mathbb{B}^{\text{red}}(\mathcal{C})$ and $\mathbb{B}_1 = \mathbb{B}^{\text{green}} = \mathbb{B}^{\text{green}}(\mathcal{C})$ by:

$\mathbb{B}^{\text{red}}$ is the Boolean algebra freely generated by $\{x_i : i < \lambda\}$ except that

$$\begin{aligned} &\text{if } i_0, \dots, i_{n-1} < \lambda, \nu = \eta_{i_0} \restriction \zeta = \eta_{i_1} \restriction \zeta = \dots = \eta_{i_{n-1}} \restriction \zeta \text{ and} \\ &\mathbb{B}_\nu \models \bigcap_{\ell < n} y_{\eta_{i_\ell} \restriction (\zeta+1)} = \mathbf{0} \\ &\text{then } \bigcap_{\ell < n} x_{i_\ell} = \mathbf{0} \end{aligned}$$

[Note: we may demand that the sequence $\langle \eta_{i_\ell}(\zeta) : \ell < n \rangle$ is strictly increasing, this will cause no difference.]

$\mathbb{B}^{\text{green}}$ is the Boolean algebra freely generated by $\{x_i : i < \lambda\}$ except that

$$\begin{aligned} &\text{if } \nu = \eta_i \restriction \zeta = \eta_j \restriction \zeta, \eta_i(\zeta) \neq \eta_j(\zeta) \text{ and} \\ &\mathbb{B}_\nu \models y_{\eta_i \restriction (\zeta+1)} \cap y_{\eta_j \restriction (\zeta+1)} \neq \mathbf{0} \\ &\text{then } x_i \cap x_j = \mathbf{0}. \end{aligned}$$

Remark 3.5 1. The equations for the green case can look strange but they have to be dual to the ones of the red case.

2. “Freely generated except ...” means that a Boolean combination is non-zero except when some (finitely many) conditions implies it. For this it is enough to look at elements of the form

$$x_{i_0}^{t_0} \cap \dots \cap x_{i_{n-1}}^{t_{n-1}}$$

where $t_\ell \in \{0, 1\}$.

3. Working in the free product $\mathbb{B}^{\text{red}} * \mathbb{B}^{\text{green}}$ we will use the same notation for elements (e.g. generators) of $\mathbb{B}^{\text{red}}$ as for elements of $\mathbb{B}^{\text{green}}$. Thus x_i may stay either for the respective generator in $\mathbb{B}^{\text{red}}$ or $\mathbb{B}^{\text{green}}$. We hope that this will not be confusing, as one can easily decide in which algebra the element is considered from the place of it (if $x \in \mathbb{B}^{\text{red}}$, $y \in \mathbb{B}^{\text{green}}$ then (x, y) will stay for the element $x \cap_{\mathbb{B}^{\text{red}} * \mathbb{B}^{\text{green}}} y \in \mathbb{B}^{\text{red}} * \mathbb{B}^{\text{green}}$). In particular we may write (x_i, x_i) for an element which could be denoted $x_i^{\text{red}} \cap x_i^{\text{green}}$.

Remark 3.6 If the pair $(\mathbb{B}^{\text{red}}, \mathbb{B}^{\text{green}})$ is a counterexample with the free product $\mathbb{B}^{\text{red}} * \mathbb{B}^{\text{green}}$ failing the λ -cc but each of the algebras satisfying that condition then each of the algebras fails the λ -Knaster condition. But $\mathbb{B}^{\text{red}}$ is supposed to have κ -cc (κ smaller than λ). This is known to restrict λ .

Proposition 3.7 Assume that $\mathcal{C} = (T, \bar{\lambda}, \bar{\eta}, \langle (\mathbb{B}_\eta, \bar{y}_\eta) : \eta \in T \rangle)$ is a (δ, μ, λ) -constructor and $J \supseteq J_\delta^{\text{bd}}$ is an ideal on δ such that

(a) $\bar{\eta} = \langle \eta_i : i \in T \rangle$ is J -cofinal for $(T, \bar{\lambda}, \bar{I})$

(b) if $X \in I_\eta^+$ then

$$(\exists \alpha, \beta \in X)(\mathbb{B}_\eta \models y_{\eta \hat{\wedge} \alpha} \cap y_{\eta \hat{\wedge} \beta} = \mathbf{0}).$$

Then the sequence $\langle x_\alpha^{\text{red}} : \alpha < \lambda \rangle$ exemplifies that $\mathbb{B}^{\text{red}}(\mathcal{C})$ fails the λ -Knaster condition.

Explanation: The above proposition is not just something in the direction of Problem 0.6. The tuple $(\mathbb{B}^{\text{red}}, \bar{x}, J_\lambda^{\text{bd}})$ is like $(\mathbb{B}_\eta, \bar{y}_\eta, I_\eta)$ but J_λ^{bd} is nicer than ideals given by previous results. Using such objects makes building examples for Problem 0.6 much easier.

PROOF It is enough to show that

for each $Y \in [\lambda]^\lambda$ one can find $\varepsilon, \zeta \in Y$ such that

$$\mathbb{B}_{\eta_\varepsilon \upharpoonright i} \models y_{\eta_\varepsilon \upharpoonright (i+1)} \cap y_{\eta_\zeta \upharpoonright (i+1)} = \mathbf{0}$$

where $i = \text{lg}(\eta_\varepsilon \wedge \eta_\zeta)$.

For this, for each $\nu \in T$ we put

$$A_\nu \stackrel{\text{def}}{=} \{\alpha < \lambda_\nu : (\exists \varepsilon \in Y)(\nu \hat{\wedge} \langle \alpha \rangle \triangleleft \eta_\varepsilon)\}.$$

Claim 3.7.1 *There is $\nu \in T$ such that $A_\nu \notin I_\nu$.*

Why? First note that by the definition of A_ν , for each $\varepsilon \in Y$ we have

$$(\forall i < \delta)(\eta_\varepsilon \hat{\wedge} \langle i \rangle \in A_{\eta_\varepsilon \upharpoonright i}).$$

Now, if we had that $A_\nu \in I_\nu$ for all $\nu \in T$ then we could apply the assumption that $\bar{\eta}$ is J -cofinal for $(T, \bar{\lambda}, \bar{I})$ to the sequence $\langle A_\nu : \nu \in T \rangle$. Thus we would find $\alpha^* < \lambda$ such that

$$\alpha^* \leq \alpha < \lambda \quad \Rightarrow \quad \{i < \delta : \eta_\alpha(i) \notin A_{\eta_\alpha \upharpoonright i}\} \in J$$

which contradicts our previous remark (remember $|Y| = \lambda$). The claim is proved.

Due to the claim we find $\nu \in T$ such that $A_\nu \notin I_\nu$. By the part (b) of our assumptions we find $\alpha, \beta \in A_\nu$ such that

$$\mathbb{B}_\nu \models y_{\nu \hat{\wedge} \langle \alpha \rangle} \cap y_{\nu \hat{\wedge} \langle \beta \rangle} = \mathbf{0}.$$

Choose $\varepsilon, \zeta \in Y$ such that $\nu \hat{\wedge} \langle \alpha \rangle \triangleleft \eta_\varepsilon$, $\nu \hat{\wedge} \langle \beta \rangle \triangleleft \eta_\zeta$ (see the definition of A_ν). Then $\nu = \eta_\varepsilon \wedge \eta_\zeta$ and

$$\mathbb{B}_\nu \models y_{\eta_\varepsilon \upharpoonright (i+1)} \cap y_{\eta_\zeta \upharpoonright (i+1)} = \mathbf{0}$$

(where $i = \text{lg}(\nu)$), finishing the proof of the proposition. ■

Lemma 3.8 *Let $\mathcal{C} = (T, \bar{\lambda}, \bar{\eta}, \langle (\mathbb{B}_\eta, \bar{y}_\eta) : \eta \in T \rangle)$ be a (δ, μ, λ) -constructor such that*

(★) *the Boolean algebras $\mathbb{B}_\eta$ satisfy the $(2^{|\delta|})^+$ -Knaster condition.*

Then the Boolean algebra $\mathbb{B}^{\text{red}}(\mathcal{C})$ satisfies the $(2^{|\delta|})^+$ -Knaster condition. In fact we may replace $(2^{|\delta|})^+$ above by any regular cardinal θ such that

$$(\forall \alpha < \theta)(|\alpha|^{|\delta|} < \theta).$$

To get that $\mathbb{B}^{\text{red}}(\mathcal{C})$ satisfies the $(2^{|\delta|})^+$ -cc it is enough if instead of (★) we assume

(★★) every free product of finitely many of the Boolean algebras $\mathbb{B}_\eta$ satisfies the $(2^{|\delta|})^+ - cc$.

Remark: 1. Usually we will have $\delta = \text{cf}(\mu)$.
 2. Later we will get more (e.g. $|\delta|^+$ -Knaster if $(T, \bar{\eta})$ is hereditarily free, see 5.12, 5.13).

PROOF Let $\theta = (2^{|\delta|})^+$ and assume (★) (the other cases have the same proofs). Suppose that $z_\varepsilon \in \mathbb{B}^{\text{red}} \setminus \{\mathbf{0}\}$ (for $\varepsilon < \theta$). We start with a series of reductions which we describe fully here but later, in similar situations, we will state what is the result of the procedure only.

Standard cleaning: Each z_ε is a Boolean combination of some generators $x_{i_0}, \dots, x_{i_{n-1}}$. But, as we want to find a subsequence with non-zero intersections, we may replace z_ε by any non-zero $z \leq z_\varepsilon$. Consequently we may assume that each z_ε is an intersection of some generators or their complements. Further, as $\text{cf}(\theta) = \theta > \aleph_0$ we may assume that the number of generators needed for this representation does not depend on ε and is equal to, say, n^* . Thus we have two functions

$$i : \theta \times n^* \longrightarrow \lambda \quad \text{and} \quad t : \theta \times n^* \longrightarrow 2$$

such that for each $\varepsilon < \theta$:

$$z_\varepsilon = \bigcap_{\ell < n^*} (x_{i(\varepsilon, \ell)})^{t(\varepsilon, \ell)}$$

and there is no repetition in $\langle i(\varepsilon, \ell) : \ell < n^* \rangle$. Moreover we may assume that $t(\varepsilon, \ell)$ does not depend on ε , i.e. $t(\varepsilon, \ell) = t(\ell)$. Due to the Δ -lemma for finite sets we may assume that $\langle \langle i(\varepsilon, \ell) : \ell < n^* \rangle : \varepsilon < \theta \rangle$ is a Δ -system of sequences, i.e.:

$$(*)_1 \quad i(\varepsilon, \ell_1) = i(\varepsilon, \ell_2) \Rightarrow \ell_1 = \ell_2 \quad \text{and}$$

$$(*)_2 \quad \text{for some } w \subseteq n^* \text{ we have}$$

$$(\exists \varepsilon_1 < \varepsilon_2 < \theta)(i(\varepsilon_1, \ell) = i(\varepsilon_2, \ell)) \text{ iff } (\forall \varepsilon_1, \varepsilon_2 < \theta)(i(\varepsilon_1, \ell) = i(\varepsilon_2, \ell)) \text{ iff } \ell \in w$$

Now note that, by the definition of the algebra $\mathbb{B}^{\text{red}}$,

$$(*)_3 \quad z_{\varepsilon_1} \cap z_{\varepsilon_2} = \mathbf{0} \quad \text{if and only if}$$

$$\bigcap \{x_{i(\varepsilon_1, \ell)}^{t(\ell)} : \ell < n^*, t(\ell) = 0\} \cap \bigcap \{x_{i(\varepsilon_2, \ell)}^{t(\ell)} : \ell < n^*, t(\ell) = 0\} = \mathbf{0}.$$

Consequently we may assume that

$$(\forall \ell < n^*)(\forall \varepsilon < \theta)(t(\ell) = 0).$$

Explanation of what we are going to do now: We want to replace the sequence $\langle z_\varepsilon : \varepsilon < \theta \rangle$ by a large subsequence such that the places of splitting between two branches used in two different z_ε 's will be uniform. Then we will be able to translate our θ -cc problem to the one on the algebras $\mathbb{B}_\eta$.

Let

$$A_\varepsilon \stackrel{\text{def}}{=} \{\nu \in {}^{\delta >} \mu : (\exists j < \varepsilon)(\exists \ell < n^*)(\nu \restriction \eta_{i(j, \ell)})\}$$

and let B_ε be the closure of A_ε :

$$B_\varepsilon \stackrel{\text{def}}{=} \{\rho \in {}^{\delta \geq} \mu : \rho \in A_\varepsilon \text{ or } \text{lg}(\rho) \text{ is a limit ordinal and } (\forall \zeta < \text{lg}(\rho))(\rho \restriction \zeta \in A_\varepsilon)\}$$

Note that $|A_\varepsilon| \leq |\varepsilon| \cdot |\delta|$ and hence $|B_\varepsilon| \leq |A_\varepsilon|^{\leq |\delta|} < \theta$. Next we define (for $\varepsilon < \theta$, $\ell < n^*$):

$$\zeta(\varepsilon, \ell) \stackrel{\text{def}}{=} \sup\{\zeta < \delta : \eta_{i(\varepsilon, \ell)} \restriction \zeta \in B_\varepsilon\}.$$

Thus $\zeta(\varepsilon, \ell) \leq \text{lg}(\eta_{i(\varepsilon, \ell)}) = \delta$. Let $\mathcal{S} = \{\varepsilon < \theta : \text{cf}(\varepsilon) > |\delta|\}$. For each $\varepsilon \in \mathcal{S}$ we necessarily have

$$\eta_{i(\varepsilon, \ell)} \restriction \zeta(\varepsilon, \ell) \in B_\varepsilon \quad \text{and} \quad B_\varepsilon = \bigcup_{\xi < \varepsilon} B_\xi$$

(remember that $\text{cf}(\varepsilon) > |\delta|$ and for limit ε we have $A_\varepsilon = \bigcup_{\xi < \varepsilon} A_\xi$) and hence

$$\eta_{i(\varepsilon, \ell)} \restriction \zeta(\varepsilon, \ell) \in B_{\xi(\varepsilon, \ell)}, \quad \text{for some } \xi(\varepsilon, \ell) < \varepsilon.$$

Let $\xi(\varepsilon) = \max\{\xi(\varepsilon, \ell) : \ell < n^*\}$. By the Fodor lemma we find $\xi^* < \theta$ such that the set

$$\mathcal{S}_1 \stackrel{\text{def}}{=} \{\varepsilon \in \mathcal{S} : \xi(\varepsilon) = \xi^*\}$$

is stationary. Thus $\eta_{i(\varepsilon, \ell)} \restriction \zeta(\varepsilon, \ell) \in B_{\xi^*}$ for each $\varepsilon \in \mathcal{S}_1$, $\ell < n^*$. Since $|B_{\xi^*}|, |\delta| < \theta$ we find $\nu_0, \dots, \nu_{n^*-1} \in B_{\xi^*}$ and $\alpha(\ell_1, \ell_2) \leq \delta$ (for $\ell_1 \leq \ell_2 < n^*$) such that the set

$$\mathcal{S}_2 \stackrel{\text{def}}{=} \{\varepsilon \in \mathcal{S}_1 : (\forall \ell < n^*)(\eta_{i(\varepsilon, \ell)} \restriction \zeta(\varepsilon, \ell) = \nu_\ell) \ \& \ (\forall \ell_1 \leq \ell_2 < n^*)(\text{lg}(\eta_{i(\varepsilon, \ell_1)}) \wedge \eta_{i(\varepsilon, \ell_2)}) = \alpha(\ell_1, \ell_2))\}$$

is stationary. Further, applying the Δ -lemma we find a set $\mathcal{S}_3 \in [\mathcal{S}_2]^\theta$ such that

$$\{(\eta_{i(\varepsilon, \ell)}(\text{lg}(\nu_\ell)) : \ell < n^*) : \varepsilon \in \mathcal{S}_3\}$$

forms a Δ -system of sequences.

For $\varepsilon \in \mathcal{S}_3$ and $\nu \in T$ denote

$$b_\nu^\varepsilon \stackrel{\text{def}}{=} \bigcap \{y_{\eta_{i(\varepsilon, \ell)} \upharpoonright (\lg(\nu)+1)} : \ell < n^*, \nu \triangleleft \eta_{i(\varepsilon, \ell)}\} \in \mathbb{B}_\nu.$$

Claim 3.8.1 *For each $\varepsilon \in \mathcal{S}_3$, $\nu \in T$ the element b_ν^ε (of the algebra $\mathbb{B}_\nu$) is non-zero.*

Why? Because of the definition of $\mathbb{B}^{\text{red}}$ and the fact that $z_\varepsilon \neq \mathbf{0}$:

$$b_\nu^\varepsilon = \mathbf{0} \quad \Rightarrow \quad \bigcap \{x_{\eta_{i(\varepsilon, \ell)}} : \ell < n^*, \nu \triangleleft \eta_{i(\varepsilon, \ell)}\} = \mathbf{0} \quad \Rightarrow \quad z_\varepsilon = \mathbf{0}.$$

Since for each $\ell < n^*$ the algebra $\mathbb{B}_{\nu_\ell}$ satisfies the θ -Knaster property we find a set $\mathcal{S}_4 \in [\mathcal{S}_3]^\theta$ such that for each $\ell < n^*$ and $\varepsilon_1, \varepsilon_2 \in \mathcal{S}_4$ we have

$$\varepsilon_1 \neq \varepsilon_2 \quad \Rightarrow \quad b_{\nu_\ell}^{\varepsilon_1} \cap b_{\nu_\ell}^{\varepsilon_2} \neq \mathbf{0} \quad \text{in } \mathbb{B}_{\nu_\ell}.$$

Now we may finish by proving the following claim.

Claim 3.8.2 *For each $\varepsilon_1, \varepsilon_2 \in \mathcal{S}_4$*

$$\mathbb{B}^{\text{red}} \models z_{\varepsilon_1} \cap z_{\varepsilon_2} \neq \mathbf{0}.$$

Why? Since $z_{\varepsilon_1} \cap z_{\varepsilon_2}$ is just the intersection of generators it is enough to show that (remember the definition of $\mathbb{B}^{\text{red}}$):

($\otimes$) for each $\varepsilon_1, \varepsilon_2 \in \mathcal{S}_4$ and for every $\nu \in T$

$$\mathbb{B}_\nu \models \bigcap \{y_{\eta_i \upharpoonright (\lg(\nu)+1)} : i \in \{i(\varepsilon_1, \ell), i(\varepsilon_2, \ell) : \ell < n^*\} \text{ and } \nu \triangleleft \eta_i\} \neq \mathbf{0}.$$

If $\nu = \nu_\ell$, $\ell < n^*$ then the intersection is $b_{\nu_\ell}^{\varepsilon_1} \cap b_{\nu_\ell}^{\varepsilon_2}$ which by the choice of the set $\mathcal{S}_4$ is not zero. So suppose that $\nu \notin \{\nu_\ell : \ell < n^*\}$. Put

$$u_\nu \stackrel{\text{def}}{=} \{i : \nu \triangleleft \eta_i \text{ and for some } \ell < n^* \text{ either } i = i(\varepsilon_1, \ell) \text{ or } i = i(\varepsilon_2, \ell)\}.$$

If

$$\{\eta_i(\lg(\nu)) : i \in u_\nu\} \subseteq \{\eta_{i(\varepsilon_2, \ell)}(\lg(\nu)) : \ell < n^* \text{ \& } \nu \triangleleft \eta_{i(\varepsilon_2, \ell)}\}$$

then we are done as $b_\nu^{\varepsilon_2} \neq \mathbf{0}$. So there is $\ell_1 < n^*$ such that $\nu \triangleleft \eta_{i(\varepsilon_1, \ell_1)}$ and

$$\eta_{i(\varepsilon_1, \ell_1)} \upharpoonright (\lg(\nu) + 1) \notin \{\eta_{i(\varepsilon_2, \ell)} \upharpoonright (\lg(\nu) + 1) : \ell < n^* \text{ \& } \nu \triangleleft \eta_{i(\varepsilon_2, \ell)}\}.$$

Similarly we may assume that there is $\ell_2 < n^*$ such that $\nu \triangleleft \eta_{i(\varepsilon_2, \ell_2)}$ and

$$\eta_{i(\varepsilon_2, \ell_2)} \restriction (\lg(\nu) + 1) \notin \{\eta_{i(\varepsilon_1, \ell)} \restriction (\lg(\nu) + 1) : \ell < n^* \text{ \& } \nu \triangleleft \eta_{i(\varepsilon_1, \ell)}\}.$$

Because of the symmetry we may assume that $\varepsilon_1 < \varepsilon_2$. Then

$$\nu = \eta_{i(\varepsilon_2, \ell_2)} \restriction (\lg(\nu) + 1) \in A_{\varepsilon_1+1} \subseteq B_{\varepsilon_2}$$

and hence $\zeta(\varepsilon_2, \ell_2) \geq \lg(\nu)$. By the choice of $\mathcal{S}_2$ (remember $\varepsilon_1, \varepsilon_2 \in \mathcal{S}_4 \subseteq \mathcal{S}_2$), we get $\nu \leq \nu_{\ell_2}$. But we have assumed that $\nu \neq \nu_{\ell_2}$, so $\nu \triangleleft \nu_{\ell_2}$. Hence (once again due to $\varepsilon_1, \varepsilon_2 \in \mathcal{S}_2$)

$$\eta_{i(\varepsilon_2, \ell_2)} \restriction (\lg(\nu) + 1) = \eta_{i(\varepsilon_1, \ell_2)} \restriction (\lg(\nu) + 1) = \nu_{\ell_2} \restriction (\lg(\nu) + 1)$$

which contradicts the choice of ℓ_2 .

The claim and so the lemma are proved. ■

Remark 3.9 We can strengthen “ θ -Knaster” in the assumption and conclusion of 3.8 in various ways. For example we may have that “intersection of any n members of the final set is non-zero”.

Definition 3.10 Let $(\mathbb{B}, \bar{y})$ be a λ -marked Boolean algebra, $\kappa \leq \lambda$. We say that

1. $(\mathbb{B}, \bar{y})$ satisfies the κ -Knaster property if $\mathbb{B}$ satisfies the definition of the κ -Knaster property (see 0.4) with restriction to subsequences of $\bar{y}$.
2. $(\mathbb{B}, \bar{y})$ is $(\kappa, \text{not } \lambda)$ -Knaster if
 - (a) the algebra $\mathbb{B}$ has the κ -Knaster property but
 - (b) the sequence $\bar{y}$ witnesses that the λ -Knaster property fails for $\mathbb{B}$.

Conclusion 3.11 Assume that μ is a strong limit singular cardinal, $\lambda = 2^\mu = \mu^+$ and $\theta = (2^{\text{cf}(\mu)})^+$.

Then there exists a λ -marked Boolean algebra $(\mathbb{B}, \bar{y})$ which is $(\theta, \text{not } \lambda)$ -Knaster.

PROOF Choose cardinals $\mu_i^0, \mu_i < \mu$ (for $i < \text{cf}(\mu)$) such that

$$(\alpha) \text{ cf}(\mu) < \mu_0^0$$

$$(\beta) \prod_{j < i} \mu_j < \mu_i^0, \mu_i = (2^{\mu_i^0})^+$$

(γ) the sequences $\langle \mu_i : i < \text{cf}(\mu) \rangle$, $\langle \mu_i^0 : i < \text{cf}(\mu) \rangle$ are increasing cofinal in μ

(possible as μ is strong limit singular). By proposition 3.1 we find μ_i -marked Boolean algebras $(\mathbb{B}_i, \bar{y}^i)$ and $(\mu_i^0)^+$ -complete ideals I_i on μ_i (for $i < \delta$) such that

(a) if $X \subseteq \mu_i$, $X \not\subseteq I_i$ then $(\exists \alpha, \beta \in X)(\mathbb{B}_i \models y_\alpha^i \cap y_\beta^i = \mathbf{0})$

(b) the algebra $\mathbb{B}_i$ has the $(2^{\text{cf}(\mu)})^+$ -Knaster property.

Let $T = \bigcup_{i < \text{cf}(\mu)} \prod_{j < i} \mu_j$ and for $\nu \in T_i$ ($i < \text{cf}(\mu)$) let $I_\nu = I_i$, $\mathbb{B}_\nu = \mathbb{B}_i$, $\bar{y}_\nu = \bar{y}^i$ and $\lambda_\nu = \mu_i$. Now we may apply proposition 2.8 to μ , $\langle \mu_i^0 : i < \text{cf}(\mu) \rangle$ and $(T, \bar{\lambda}, \bar{I})$ to find a stronger $J_{\text{cf}(\mu)}^{\text{bd}}$ -cofinal sequence $\bar{\eta}$ for $(T, \bar{\lambda}, \bar{I})$ of the length λ . Consider the $(\text{cf}(\mu), \mu, \lambda)$ -constructor $\mathcal{C} = (T, \bar{\lambda}, \bar{\eta}, \langle (\mathbb{B}_\nu, \bar{y}_\nu) : \nu \in T \rangle)$. By (b) above we may apply lemma 3.8 to get that the algebra $\mathbb{B}^{\text{red}}(\mathcal{C})$ satisfies the $(2^{\text{cf}(\mu)})^+$ -Knaster condition. Finally we use proposition 3.7 (and (a) above) to conclude that $(\mathbb{B}^{\text{red}}(\mathcal{C}), \langle x_\alpha^{\text{red}} : \alpha < \lambda \rangle)$ is $(\theta, \text{not } \lambda)$ -Knaster. ■

Proposition 3.12 *Assume that:*

κ is a regular cardinal such that $(\forall \alpha < \kappa)(|\alpha|^{|\delta|} < \kappa)$, $\bar{\lambda} = \langle \lambda_i : i < \delta \rangle$ is an increasing sequence of regular cardinals such that $\kappa \leq \lambda_0$, $\prod_{j < i} \lambda_j < \lambda_i$ (or just $\max \text{pcf}\{\lambda_j : j < i\} < \lambda_i$) for $i < \delta$ and $\lambda \in \text{pcf}\{\lambda_i : i < \delta\}$. Further suppose that for each $i < \delta$ there exists a λ_i -marked Boolean algebra which is $(\kappa, \text{not } \lambda_i)$ -Knaster.

Then there exists a λ -marked Boolean algebra which is $(\kappa, \text{not } \lambda)$ -Knaster.

PROOF If $\lambda = \lambda_i$ for some $i < \delta$ then there is nothing to do. If $\lambda < \lambda_i$ for some $i < \delta$ then let $\alpha < \delta$ be the maximal limit ordinal such that $(\forall i < \alpha)(\lambda_i < \lambda)$ (it necessarily exists). Now we may replace $\langle \lambda_i : i < \delta \rangle$ by $\langle \lambda_i : i < \alpha \rangle$. Thus we may assume that $(\forall i < \delta)(\lambda_i < \lambda)$. Further we may assume that

$$\lambda = \max \text{pcf}\{\lambda_i : i < \delta\}$$

(by [Sh:g], I, 1.8). Now, due to [Sh:g], II, 3.5, p.65, we find a sequence $\bar{\eta} \subseteq \prod_{i < \delta} \lambda_i$ and an ideal J on δ such that

1. $J \supseteq J_\delta^{\text{bd}}$ and $\lambda = \text{tcf}(\prod_{i < \delta} \lambda_i / J)$
(naturally: $J = \{a \subseteq \delta : \max \text{pcf}\{\lambda_i : i \in a\} < \lambda\}$)
2. $\bar{\eta} = \langle \eta_\varepsilon : \varepsilon < \lambda \rangle$ is $<_J$ -increasing cofinal in $\prod_{i < \delta} \lambda_i / J$

3. for each $i < \delta$

$$|\{\eta_\varepsilon \restriction i : \varepsilon < \lambda\}| < \lambda_i.$$

Let $T = \bigcup_{i < \delta} \prod_{j < i} \lambda_j$ and for $\nu \in T_i$ ($i < \delta$) let $\lambda_\nu = \lambda_i$, $I_\nu = J_{\lambda_i}^{\text{bd}}$.

It follows from the choice of $\bar{\eta}, J$ above and our assumptions that we may apply proposition 2.6 and hence $\bar{\eta}$ is J -cofinal for $(T, \bar{\lambda}, \bar{I})$. For $\nu \in T$ let $(\mathbb{B}_\nu, \bar{y}_\nu)$ be a λ_ν -marked $(\kappa, \text{not } \lambda_\nu)$ -Knaster Boolean algebra (exists by our assumptions). Now we may finish using 3.8 and 3.7 for $\mathcal{C} = (T, \bar{\lambda}, \bar{\eta}, \langle (\mathbb{B}_\eta, \bar{y}_\eta) : \eta \in T \rangle)$, $\bar{I}$ and J (note the assumption **(b)** of 3.7 is satisfied as $I_\eta = J_{\lambda_\eta}^{\text{bd}}$; remember the choice of $(\mathbb{B}_\eta, \bar{y}_\eta)$). ■

Remark 3.13 Note that from cardinal arithmetic hypothesis $\text{cf}(\mu) = \chi$, $\chi^{<\chi} < \chi < \mu$, $\mu^+ = \lambda < 2^\chi$ alone we cannot hope to build a counterexample. This is because of section 4 of [Sh:93], particularly lemma 4.13 there. It was shown in that paper that if $\chi^{<\chi} < \chi_1 = \chi_1^{<\chi_1}$ then there is a χ^+ -cc χ -complete forcing notion $\mathbb{P}$ of size χ_1 such that

$$\begin{aligned} \Vdash_{\mathbb{P}} \quad & \text{“if } |\mathbb{B}| < \chi_1, \mathbb{B} \models \chi\text{-cc} \\ & \text{then } \mathbb{B}^+ \text{ is the union of } \chi \text{ ultrafilters”}. \end{aligned}$$

More on this see in section 8.

So the centrality of $\lambda \in \text{Reg} \cap (\mu, 2^\mu]$, μ strong limit singular, is very natural.

4 The main result

Proposition 4.1 *Suppose that $\mathcal{C}$ is a (δ, μ, λ) -constructor. Then the free product $\mathbb{B}^{\text{red}}(\mathcal{C}) * \mathbb{B}^{\text{green}}(\mathcal{C})$ fails the λ -cc (so $\text{c}(\mathbb{B}^{\text{red}}(\mathcal{C}) * \mathbb{B}^{\text{green}}(\mathcal{C})) \geq \lambda$).*

PROOF Look at the elements $(x_i, x_i) \in \mathbb{B}^{\text{red}} * \mathbb{B}^{\text{green}}$ for $i < \lambda$. It follows directly from the definition of the algebras that for each $i < j < \lambda$:

$$\text{either } \mathbb{B}^{\text{red}} \models x_i^{\text{red}} \cap x_j^{\text{red}} = \mathbf{0} \quad \text{or} \quad \mathbb{B}^{\text{green}} \models x_i^{\text{green}} \cap x_j^{\text{green}} = \mathbf{0}.$$

Consequently the sequence $\langle (x_i, x_i) : i < \lambda \rangle$ witnesses the assertion of the proposition. ■

Proposition 4.2 *Suppose that $n < \omega$ and for $\ell \leq n$:*

1. χ_ℓ, λ_ℓ are regular cardinals, $\chi_\ell < \lambda_\ell < \chi_{\ell+1}$
2. $(\mathbb{B}_\ell, \bar{y}_\ell, I_\ell)$ is a $(\lambda_\ell, \chi_\ell)$ -well marked Boolean algebra (see definition 3.2), $\bar{y}_\ell = \langle y_i^\ell : i < \lambda_\ell \rangle$

3. $\mathbb{B}$ is the Boolean algebra freely generated by $\{y_\eta : \eta \in \prod_{\ell \leq n} \lambda_\ell\}$ except that

$$\begin{aligned} & \text{if } \eta_{i_0}, \dots, \eta_{i_{k-1}} \in \prod_{\ell \leq n} \lambda_\ell, \eta_{i_0} \restriction \ell = \eta_{i_1} \restriction \ell = \dots = \eta_{i_{n-1}} \restriction \ell \text{ and} \\ & \mathbb{B}_\ell \models \bigcap_{m < k} y_{\eta_{i_m}(\ell)}^\ell = \mathbf{0} \\ & \text{then } \bigcap_{m < k} y_{\eta_{i_m}} = \mathbf{0}. \end{aligned}$$

[Compare to the definition of the algebras $\mathbb{B}^{\text{red}}(\mathcal{C})$.]

4. $I = \{B \subseteq \prod_{\ell \leq n} \lambda_\ell : \neg(\exists^{I_0} \gamma_0) \dots (\exists^{I_n} \gamma_n)(\langle \gamma_0, \dots, \gamma_n \rangle \in B)\}$.

Then:

- (a) if all the algebras $\mathbb{B}_\ell$ (for $\ell \leq n$) satisfy the θ -Knaster property, θ is a regular uncountable cardinal then $\mathbb{B}$ has the θ -Knaster property;
- (b) I is a χ_0 -complete ideal on $\prod_{\ell \leq n} \lambda_i$;
- (c) if $Y \subseteq (\prod_{\ell \leq n} \lambda_\ell)^n$ is such that

$$(\exists^I \eta_0) \dots (\exists^I \eta_n)(\langle \eta_0, \dots, \eta_n \rangle \in Y)$$

then there are $\langle \eta'_0, \dots, \eta'_n \rangle, \langle \eta''_0, \dots, \eta''_n \rangle \in Y$ such that for all $\ell \leq n$

$$\mathbb{B} \models y_{\eta'_\ell} \cap y_{\eta''_\ell} = \mathbf{0}.$$

PROOF (a) The proof that the algebra $\mathbb{B}$ satisfies θ -Knaster condition is exactly the same as that of 3.8 (actually it is a special case of that).

(b) Should be clear.

(c) For $\ell, m \leq n$ put

$$\chi_\ell^m = \chi_\ell, \lambda_\ell^m = \lambda_\ell, I_\ell^m = I_\ell, P_\ell^m = \{\{\alpha, \beta\} \subseteq \lambda_\ell : \mathbb{B}_\ell \models y_\alpha^\ell \cap y_\beta^\ell = \mathbf{0}\}, B = Y.$$

It is easy to check that the assumptions of proposition 1.5 are satisfied. Applying it we find sets $X_0, \dots, X_n$ satisfying the respective versions of clauses (a)–(d) there. Note that our choice of the sets P_ℓ^m and clauses (b), (c) of 1.5 imply that

$$\begin{aligned} X_m &= \{\nu'_m, \nu''_m\} \subseteq \prod_{\ell \leq n-m} \lambda_\ell \\ \nu'_m \restriction (n-m) &= \nu''_m \restriction (n-m) \\ \mathbb{B}_{n-m} &\models y_{\nu'_m(n-m)}^{n-m} \cap y_{\nu''_m(n-m)}^{n-m} = \mathbf{0} \end{aligned}$$

Look at the sequences $\langle \nu'_0, \dots, \nu'_n \rangle, \langle \nu''_0, \dots, \nu''_n \rangle$. By the clause **(d)** of 1.5 we find $\langle \eta'_0, \dots, \eta'_m \rangle, \langle \nu''_0, \dots, \nu''_n \rangle \in Y$ such that for each $m \leq n$

$$\nu'_m \trianglelefteq \eta'_m, \quad \nu''_m \trianglelefteq \eta''_m.$$

Now, the properties of ν'_m, ν''_m and the definition of the algebra $\mathbb{B}$ imply that for each $m \leq n$:

$$\mathbb{B} \models y_{\eta'_m} \cap y_{\eta''_m} = \mathbf{0},$$

finishing the proof. ■

Lemma 4.3 *Assume that λ is a regular cardinal, $|\delta| < \lambda$, J is an ideal on δ extending J_δ^{bd} , $\mathcal{C} = (T, \bar{\lambda}, \bar{\eta}, \langle (\mathbb{B}_\eta, \bar{y}_\eta) : \eta \in T \rangle)$ is a (δ, μ, λ) -constructor and $\bar{I}$ is such that $(T, \bar{\lambda}, \bar{I}) \in \mathcal{K}_{\delta, \mu}^{\text{id}}$. Suppose that $\bar{\eta} = \langle \eta_\alpha : \alpha < \lambda \rangle$ is a stronger (or big) J -cofinal in $(T, \bar{\lambda}, \bar{I})$ sequence such that*

$$(\forall i < \delta)(|\{\eta_\alpha \restriction i : \alpha < \lambda\}| < \lambda).$$

Further assume that

($\ominus$) for every $n < \omega$ for a J -positive set of $i < \delta$ we have:

if $\eta_0, \dots, \eta_n \in T_i$ are pairwise distinct and the set $Y \subseteq \prod_{\ell \leq n} \lambda_{\eta_\ell}$ is such that

$$(\exists^{I_{\eta_0}} \gamma_0) \dots (\exists^{I_{\eta_n}} \gamma_n) (\langle \gamma_0, \dots, \gamma_n \rangle \in Y)$$

then for some $\gamma'_\ell, \gamma''_\ell < \lambda_{\eta_\ell}$ (for $\ell \leq n$) we have

$$\langle \gamma'_\ell : \ell \leq n \rangle, \langle \gamma''_\ell : \ell \leq n \rangle \in Y \quad \text{and for all } \ell \leq n$$

$$\mathbb{B}_{\eta_\ell} \models y_{\eta_\ell \restriction \langle \gamma'_\ell \rangle} \cap y_{\eta_\ell \restriction \langle \gamma''_\ell \rangle} = \mathbf{0}.$$

Then the Boolean algebra $\mathbb{B}^{\text{green}}(\mathcal{C})$ satisfies λ -cc.

PROOF Suppose that $\langle z_\alpha : \alpha < \lambda \rangle \subseteq \mathbb{B}^{\text{green}} \setminus \{\mathbf{0}\}$. By the standard cleaning (compare the first part of the proof of 3.8) we may assume that there are $n^* \in \omega$ and a function $\varepsilon : \lambda \times n^* \longrightarrow \lambda$ such that

1. $z_\alpha = \bigcap_{\ell < n^*} x_{\varepsilon(\alpha, \ell)}$ (in $\mathbb{B}^{\text{green}}$)
2. $\varepsilon(\alpha, 0) < \varepsilon(\alpha, 1) < \dots < \varepsilon(\alpha, n^* - 1)$

3. $\langle \langle \varepsilon(\alpha, \ell) : \ell < n^* \rangle : \alpha < \lambda \rangle$ forms a Δ -system of sequences with the kernel m^* , i.e. $(\forall \ell < m^*)(\varepsilon(\alpha, \ell) = \varepsilon(\ell))$ and

$$(\forall \ell \in [m^*, n^*])(\forall \alpha < \lambda)(\varepsilon(\alpha, \ell) \notin \{\varepsilon(\beta, k) : (\beta, k) \neq (\alpha, \ell)\})$$

4. there is $i^* < \delta$ such that for each $\alpha < \lambda$ there is no repetition in the sequence $\langle \eta_{\varepsilon(\alpha, \ell)} \upharpoonright i^* : \ell < n^* \rangle$.

Since $|\{\eta_\alpha \upharpoonright i : \alpha < \lambda\}| < \lambda$ (for $i < \delta$) and $|\delta| < \lambda$ we may additionally require that

($\hat{*}$) for each $i < \delta$, for every $\alpha < \lambda$ we have

$$(\exists^\lambda \beta < \lambda)(\forall \ell < n^*)(\eta_{\varepsilon(\alpha, \ell)} \upharpoonright (i+1) = \eta_{\varepsilon(\beta, \ell)} \upharpoonright (i+1))$$

and

($\hat{**}$) for each $\alpha < \beta < \lambda$, $\ell < n^*$

$$\eta_{\varepsilon(\alpha, \ell)} \upharpoonright i^* = \eta_{\varepsilon(\beta, \ell)} \upharpoonright i^*.$$

Remark: Note that the claim below is like an $(n^* - m^*)$ -place version of 3.7. Having an $(n^* - m^*)$ -ary version is extra for the construction but it also costs.

Claim 4.3.1 Assume that:

$\mathcal{C} = (T, \bar{\lambda}, \bar{\eta}, \langle (\mathbb{B}_\eta, \bar{y}_\eta) : \eta \in T \rangle)$ is a (δ, μ, λ) -constructor, λ a regular cardinal, $\delta < \lambda$, $\bar{I}$ is such that $(T, \bar{\lambda}, \bar{I}) \in \mathcal{K}_{\delta, \mu}^{\text{id}}$, J is an ideal on δ extending J_δ^{bd} and the sequence $\bar{\eta}$ is stronger J -cofinal in $(T, \bar{\lambda}, \bar{I})$.

Further suppose that $\varepsilon : \lambda \times n^* \longrightarrow m^*, n^*$ and $i^* < \delta$ are as above (after the reduction, but the property ($\hat{**}$) is not needed).

Then

($\boxtimes$) for every large enough $\alpha < \lambda$ the set:

$$Z_\alpha \stackrel{\text{def}}{=} \{i < \delta : \neg(\exists^{I_{\eta_{\varepsilon(\alpha, m^*)}} \upharpoonright i} \gamma_{m^*})(\exists^{I_{\eta_{\varepsilon(\alpha, m^*+1)}} \upharpoonright i} \gamma_{m^*+1}) \dots \\ \dots (\exists^{I_{\eta_{\varepsilon(\alpha, n^*-1)}} \upharpoonright i} \gamma_{n^*-1})(\exists^\lambda \beta)(\forall \ell \in [m^*, n^*])(\eta_{\varepsilon(\beta, \ell)} \upharpoonright (i+1) = (\eta_{\varepsilon(\alpha, \ell)} \upharpoonright i)^\wedge \langle \gamma_\ell \rangle)\}$$

is in the ideal J .

Why? For each $i < \delta$, $i \geq i^*$ and distinct sequences $\nu_{m^*}, \dots, \nu_{n^*-1} \in T_i$ define

$$B_{\langle \nu_\ell : \ell \in [m^*, n^*] \rangle} \stackrel{\text{def}}{=} \{\bar{\gamma} : \bar{\gamma} = \langle \gamma_\ell : \ell \in [m^*, n^*] \rangle \text{ and} \\ \text{for arbitrarily large } \alpha < \lambda \text{ for all } m^* \leq \ell < n^* \\ \nu_\ell^\wedge \langle \gamma_\ell \rangle \triangleleft \eta_{\varepsilon(\alpha, \ell)}\}.$$

We will call a sequence $\langle \nu_\ell : \ell \in [m^*, n^*] \rangle$ a *success* if

$$(\exists^{I_{\nu_{m^*}} \gamma_{m^*}}) \dots (\exists^{I_{\nu_{n^*-1}} \gamma_{n^*-1}})(\langle \gamma_\ell : \ell \in [m^*, n^*] \rangle \in B_{\langle \nu_\ell : \ell \in [m^*, n^*] \rangle}).$$

Using this notion we may reformulate $(\boxtimes)$ (which we have to prove) to

$(\boxtimes^*)$ for every large enough $\alpha < \lambda$, for J -majority of $i < \delta$, $i > i^*$ the sequence $\langle \eta_{\varepsilon(\alpha, \ell)} \upharpoonright i : \ell \in [m^*, n^*] \rangle$ is a success.

To show $(\boxtimes^*)$ note that if a sequence $\langle \nu_\ell : \ell \in [m^*, n^*] \rangle$ is not a success then there are functions $f_{\langle \nu_\ell : \ell \in [m^*, n^*] \rangle}^k$ (for $m^* \leq k < n^*$) such that

$$f_{\langle \nu_\ell : \ell \in [m^*, n^*] \rangle}^k : \prod_{\ell=m^*}^{k-1} \lambda_{\nu_\ell} \longrightarrow I_{\nu_k} \quad \text{and}$$

$$\begin{aligned} \text{if} \quad & \langle \gamma_\ell : \ell \in [m^*, n^*] \rangle \in B_{\langle \nu_\ell : \ell \in [m^*, n^*] \rangle} \\ \text{then} \quad & (\exists k \in [m^*, n^*])(\gamma_k \in f_{\langle \nu_\ell : \ell \in [m^*, n^*] \rangle}^k(\gamma_{m^*}, \dots, \gamma_{k-1})). \end{aligned}$$

If $\langle \nu_\ell : \ell \in [m^*, n^*] \rangle$ is a success then we declare that $f_{\langle \nu_\ell : \ell \in [m^*, n^*] \rangle}^k$ is constantly equal to $\emptyset$.

Now we may finish the proof of the claim applying clause **(b)** of definition 2.2(5) to $n^* - 1$ and functions $F_0, \dots, F_{n^*-1}$ such that for $k \in [m^*, n^*]$

$$F_k(\nu_0 \hat{\ } \langle \gamma_0 \rangle, \dots, \nu_{k-1} \hat{\ } \langle \gamma_{k-1} \rangle, \nu_k, \dots, \nu_{n^*-1}) = f_{\langle \nu_\ell : \ell \in [m^*, n^*] \rangle}^k(\gamma_{m^*}, \dots, \gamma_{k-1}).$$

This gives us a suitable $\alpha^* < \lambda$. Suppose $\varepsilon(\alpha, m^*) \geq \alpha^*$. Then for J -majority of $i < \delta$ for each $k \in [m^*, n^*]$ we have

if

$$F_m(\eta_{\varepsilon(\alpha, 0)} \upharpoonright (i+1), \dots, \eta_{\varepsilon(\alpha, k-1)} \upharpoonright (i+1), \eta_{\varepsilon(\alpha, k)} \upharpoonright i, \dots, \eta_{\varepsilon(\alpha, n^*-1)} \upharpoonright i) \in I_{\eta_{\varepsilon(\alpha, k)} \upharpoonright i}$$

then

$$\eta_{\varepsilon(\alpha, k)} \upharpoonright (i+1) \notin F_m(\eta_{\varepsilon(\alpha, 0)} \upharpoonright (i+1), \dots, \eta_{\varepsilon(\alpha, k-1)} \upharpoonright (i+1), \eta_{\varepsilon(\alpha, k)} \upharpoonright i, \dots, \eta_{\varepsilon(\alpha, n^*-1)} \upharpoonright i).$$

But the choice of the functions F_k implies that thus for J -majority of $i < \delta$, for each $k \in [m^*, n^*]$

$$\eta_{\varepsilon(\alpha, k)}(i) \notin f_{\langle \eta_{\varepsilon(\alpha, \ell)} \upharpoonright i : \ell \in [m^*, n^*] \rangle}^k(\eta_{\varepsilon(\alpha, m^*)}(i), \dots, \eta_{\varepsilon(\alpha, k-1)}(i)).$$

Now the definition of the function $f_{\langle \nu_\ell : \ell \in [m^*, n^*] \rangle}^k$ works:

if for some relevant $i < \delta$ above the sequence $\langle \eta_{\varepsilon(\alpha, \ell)} \upharpoonright i : \ell \in [m^*, n^*] \rangle$ is not a success then $\langle \eta_{\varepsilon(\alpha, \ell)}(i) : \ell \in [m^*, n^*] \rangle \notin B_{\langle \eta_{\varepsilon(\alpha, \ell)} \upharpoonright i : \ell \in [m^*, n^*] \rangle}$ and this contradicts $\hat{*}$ before.

The claim is proved.

Let α^* be such that for each $\alpha \geq \alpha^*$ we have $Z_\alpha \in J$. Choose $i \in \delta \setminus Z_{\alpha^*}$ such that the clause $(\ominus)$ applies for $n^* - m^*$ and i . Let

$$Y \stackrel{\text{def}}{=} \{ \langle \gamma_{m^*}, \dots, \gamma_{n^*-1} \rangle : (\exists^\lambda \beta) (\forall \ell \in [m^*, n^*]) (\eta_{\varepsilon(\beta, \ell)} \upharpoonright (i+1) = (\eta_{\varepsilon(\alpha^*, \ell)} \upharpoonright i)^\wedge \langle \gamma_\ell \rangle) \}.$$

The definition of Z_{α^*} (and the choice of i) imply that the assumption $(\ominus)$ applies to the set Y and we get $\gamma'_\ell, \gamma''_\ell < \lambda_{\eta_{\varepsilon(\alpha^*, \ell)} \upharpoonright i}$ (for $m^* \leq \ell < n^*$) such that

$$\langle \gamma'_\ell : m^* \leq \ell < n^* \rangle, \langle \gamma''_\ell : m^* \leq \ell < n^* \rangle \in Y \quad \text{and}$$

$$\mathbb{B}_{\eta_{\varepsilon(\alpha^*, \ell)} \upharpoonright i} \models y_{\eta_{\varepsilon(\alpha^*, \ell)} \upharpoonright i} \langle \gamma'_\ell \rangle \cap y_{\eta_{\varepsilon(\alpha^*, \ell)} \upharpoonright i} \langle \gamma''_\ell \rangle = \mathbf{0} \quad \text{for } m^* \leq \ell < n^*.$$

Now choose $\alpha < \beta < \lambda$ such that for $m^* \leq \ell < n^*$

$$\eta_{\varepsilon(\alpha^*, \ell)} \upharpoonright i^\wedge \langle \gamma'_\ell \rangle = \eta_{\varepsilon(\alpha, \ell)} \upharpoonright (i+1), \quad \eta_{\varepsilon(\alpha^*, \ell)} \upharpoonright i^\wedge \langle \gamma''_\ell \rangle = \eta_{\varepsilon(\beta, \ell)} \upharpoonright (i+1)$$

(possible by the choice of Y and $\gamma'_\ell, \gamma''_\ell$). The definition of the algebra $\mathbb{B}^{\text{green}}(\mathcal{C})$ and the choice of $\gamma'_\ell, \gamma''_\ell$ imply that for $m^* \leq \ell < n^*$

$$\mathbb{B}^{\text{green}}(\mathcal{C}) \models x_{\varepsilon(\alpha, \ell)} \cap x_{\varepsilon(\beta, \ell)} \neq \mathbf{0}.$$

If $\ell \neq m$ then

$$\mathbb{B}^{\text{green}}(\mathcal{C}) \models x_{\varepsilon(\alpha, \ell)} \cap x_{\varepsilon(\beta, m)} \neq \mathbf{0}$$

by the conditions $(\hat{**})$ and 4) of the preliminary cleaning (and the definition of $\mathbb{B}^{\text{green}}(\mathcal{C})$, remember $z_\alpha \neq \mathbf{0}$). Finally, remembering that $\varepsilon(\alpha, \ell) = \varepsilon(\beta, \ell)$ for $\ell < m^*$, $z_\alpha \neq \mathbf{0}$ and $z_\beta \neq \mathbf{0}$, we may conclude that

$$\mathbb{B}^{\text{green}}(\mathcal{C}) \models \bigcap_{\ell < n^*} x_{\varepsilon(\alpha, \ell)} \cap \bigcap_{\ell < n^*} x_{\varepsilon(\beta, \ell)} \neq \mathbf{0}$$

finishing the proof. ■

Theorem 4.4 If μ is a strong limit singular cardinal, $\lambda \stackrel{\text{def}}{=} 2^\mu = \mu^+$ then there are Boolean algebras $\mathbb{B}_1, \mathbb{B}_2$ such that the algebra $\mathbb{B}_1$ satisfies the λ -cc, the algebra $\mathbb{B}_2$ has the $(2^{\text{cf}(\mu)})^+$ -Knaster property but the free product $\mathbb{B}_1 * \mathbb{B}_2$ does not satisfy the λ -cc.

PROOF Let $\delta = \text{cf}(\mu)$ and let $h : \delta \longrightarrow \omega$ be a function such that

$$(\forall n \in \omega)(\exists^\delta i)(h(i) = n).$$

Choose an increasing sequence $\langle \mu_i : i < \delta \rangle$ of regular cardinals such that $\mu = \sum_{i < \delta} \mu_i$. Next, by induction on $i < \delta$ choose λ_i , χ_i , $(\mathbb{B}_i, \bar{y}_i)$ and I_i such that

1. λ_i, χ_i are regular cardinals below μ
2. $\lambda_i > \chi_i \geq \prod_{j < i} \lambda_j + \mu_i$
3. I_i is a χ_i^+ -complete ideal on λ_i (containing all singletons)
4. $(\mathbb{B}_i, \bar{y}_i)$ is a λ_i -marked Boolean algebra such that
if $n = h(i)$ and the set $Y \subseteq (\lambda_i)^{n+1}$ is such that

$$(\exists^{I_i} \gamma_0) \dots (\exists^{I_i} \gamma_n) (\langle \gamma_0, \dots, \gamma_n \rangle \in Y)$$

then for some $\gamma'_\ell, \gamma''_\ell < \lambda_i$ (for $\ell \leq n$) we have

$$\langle \gamma'_\ell : \ell \leq n \rangle, \langle \gamma''_\ell : \ell \leq n \rangle \in Y \quad \text{and for all } \ell \leq n$$

$$\mathbb{B}_i \models y_{\gamma'_\ell}^i \cap y_{\gamma''_\ell}^i = \mathbf{0}.$$

5. Each algebra $\mathbb{B}_i$ satisfies the $(2^{|\delta|})^+$ -Knaster condition.

Arriving at the stage i of the construction first we put $\chi_i = (\prod_{j < i} \lambda_j + \mu_i)^+$.

Next we define inductively $\chi_{i,k}, \lambda_{i,k}$ for $k \leq h(i)$ such that

$$\chi_{i,0} = \chi_i, \quad \lambda_{i,k} = (2^{\chi_{i,k}})^+, \quad \chi_{i,k+1} = (\lambda_{i,k})^+.$$

By 3.1, for each $k \leq h(i)$ we find a $(\lambda_{i,k}, \chi_{i,k}^+)$ -well marked Boolean algebra $(\mathbb{B}_{i,k}, \bar{y}_{i,k}, I_{i,k})$ such that $\mathbb{B}_{i,k}$ has the $(2^\delta)^+$ -Knaster property (compare 3.3). Let $\lambda_i = \lambda_{i,h(i)}$. Proposition 4.2 applied to $\langle (\mathbb{B}_{i,k}, \bar{y}_{i,k}, I_{i,k}) : k \leq h(i) \rangle$ provides a λ_i -marked Boolean algebra $(\mathbb{B}_i, \bar{y}_i)$ and a χ_i^+ -complete ideal I_i on λ_i such that the requirements 4,5 above are satisfied.

Now put $T = \bigcup_{j < \delta} \prod_{i < j} \lambda_i$ and for $\eta \in T$:

$$\mathbb{B}_\eta = \mathbb{B}_{\text{lg}(\eta)}, \quad \bar{y}_\eta = \bar{y}_{\text{lg}(\eta)}, \quad I_\eta = I_{\text{lg}(\eta)}.$$

By 2.8 we find a stronger J_δ^{bd} -cofinal sequence $\bar{\eta} = \langle \eta_\alpha : \alpha < \lambda \rangle$ for $(T, \bar{\lambda}, \bar{I})$. Take the (δ, μ, μ^+) -constructor $\mathcal{C}$ determined by these parameters. Look at

the algebras $\mathbb{B}_2 = \mathbb{B}^{\text{red}}(\mathcal{C})$, $\mathbb{B}_1 = \mathbb{B}^{\text{green}}(\mathcal{C})$. Applying 4.1 we get that $\mathbb{B}_1 * \mathbb{B}_2$ fails the λ -cc. The choice of the function h and the requirement 4 above allow us to apply 4.3 to conclude that the algebra $\mathbb{B}_2$ satisfies λ -cc. Finally, by 3.8, we have that $\mathbb{B}_1$ has the $(2^\delta)^+$ -Knaster property. ■

Remark 4.5 1. We shall later give results not using $2^\mu = \mu^+$ but still not in ZFC

2. Applying the methods of [Sh 576] one can the consistency of: *for some μ strong limit singular there is no example for $\lambda = \mu^+$.*
3. If we want “for no regular $\lambda \in [\mu, 2^\mu]$ ” more is needed, we expect the consistency, but it is harder (not speaking of “for all μ ”)
4. Remark 1) above shows that $2^\mu > \mu^+$ is not enough for the negative result.

5 Toward improvements

Definition 5.1 Let $(T, \bar{\lambda}, \bar{I}) \in \mathcal{K}_{\mu, \delta}^{\text{id}}$ and let J be an ideal on δ (including J_δ^{bd} , as usual). We say that a sequence $\bar{\eta} = \langle \eta_\alpha : \alpha < \lambda \rangle$ of δ -branches through T is super J -cofinal for $(T, \bar{\lambda}, \bar{I})$ if

- (a) $\eta_\alpha \neq \eta_\beta$ for distinct $\alpha, \beta < \lambda$
- (b) for every function F there is $\alpha^* < \lambda$ such that
if $\alpha_0 < \dots < \alpha_n < \lambda$, $\alpha^* \leq \alpha_n$
then the set

$$\begin{aligned} \{i < \delta : & \text{ (ii)* } F(\eta_{\alpha_0}, \dots, \eta_{\alpha_{n-1}}, \eta_{\alpha_n} \restriction i) \in I_{\eta_{\alpha_n} \restriction i} \\ & \text{ (and well defined) but} \\ & \eta_{\alpha_n} \restriction (i+1) \in F(\eta_{\alpha_0}, \dots, \eta_{\alpha_{n-1}}, \eta_{\alpha_n} \restriction i)\} \end{aligned}$$

is in the ideal J .

Remark 5.2 1. The main difference between the definition of super J -cofinal sequence and those in 2.2 is the fact that here the values of the function F depend on η_{α_ℓ} (for $\ell < n$), not on the restrictions of these sequences as it was in earlier notions.

2. “super* J -cofinal” is defined by adding “ $\alpha^* \leq \alpha_0$ ” (compare 2.2(10)).

Proposition 5.3 *Suppose that $(T, \bar{\lambda}, \bar{I}) \in \mathcal{K}_{\mu, \delta}^{\text{id}}$ is such that for each $\nu \in T_i$, $i < \delta$ the ideal I_ν is $|T_i|^+$ -complete. Let $J \supseteq J_\delta^{\text{bd}}$ be an ideal on δ . Then every super J -cofinal sequence is stronger* J -cofinal.*

PROOF Assume that $\bar{\eta} = \langle \eta_\alpha : \alpha < \lambda \rangle \subseteq \lim_\delta(T)$ is super J -cofinal for $(T, \bar{\lambda}, \bar{I})$. Let $n < \omega$ and let $F_0, \dots, F_{n-1}$ be functions. For each $\ell \leq n$ we define an $(\ell + 1)$ -place function F_ℓ^* such that

if $\alpha_0 < \alpha_1 < \dots < \alpha_{\ell-1} < \lambda$, $\rho \in T_i$, $i < \delta$
then

$$F_\ell^*(\eta_{\alpha_0}, \dots, \eta_{\alpha_{\ell-1}}, \rho) = \bigcup \{ F_\ell(\eta_{\alpha_0} \upharpoonright (i+1), \dots, \eta_{\alpha_{\ell-1}} \upharpoonright (i+1), \rho, \nu_{\ell+1}, \dots, \nu_n) : \\ \nu_{\ell+1}, \dots, \nu_n \in T_i \text{ \& } F_\ell(\eta_{\alpha_0} \upharpoonright (i+1), \dots, \eta_{\alpha_{\ell-1}} \upharpoonright (i+1), \rho, \nu_{\ell+1}, \dots, \nu_n) \in I_\rho \\ \text{(and well defined)} \}.$$

As the ideals I_ρ (for $\rho \in T_i$) are $|T_i|^+$ -complete we know that

$$F_\ell^*(\eta_{\alpha_0}, \dots, \eta_{\alpha_{\ell-1}}, \rho) \in I_\rho.$$

Applying 5.1(b) to the functions F_ℓ^* ($\ell < n$) we choose $\alpha_\ell^* < \lambda$ such that

if $\alpha_0 < \dots < \alpha_\ell < \lambda$, $\alpha_\ell^* \leq \alpha_\ell$
then the set

$$B_\ell^* \stackrel{\text{def}}{=} \{ i < \delta : F_\ell^*(\eta_{\alpha_0}, \dots, \eta_{\alpha_{\ell-1}}, \eta_{\alpha_\ell} \upharpoonright i) \in I_{\eta_{\alpha_\ell} \upharpoonright i} \text{ but } \\ \eta_{\alpha_\ell} \upharpoonright (i+1) \in F_\ell^*(\eta_{\alpha_0}, \dots, \eta_{\alpha_{\ell-1}}, \eta_{\alpha_\ell} \upharpoonright i) \}$$

is in the ideal J .

Put $\alpha^* = \max\{\alpha_\ell^* : \ell \leq n\}$. We want to show that this α^* works for the condition 2.2(6)(b) (version for “stronger*”). So suppose that $m \leq n$, $\alpha^* \leq \alpha_0 < \alpha_1 < \dots < \alpha_n < \lambda$. Let

$$B_m \stackrel{\text{def}}{=} \{ i < \delta : F_m(\eta_{\alpha_0} \upharpoonright (i+1), \dots, \eta_{\alpha_{m-1}} \upharpoonright (i+1), \eta_{\alpha_m} \upharpoonright i, \dots, \eta_{\alpha_n} \upharpoonright i) \in I_{\eta_{\alpha_m} \upharpoonright i} \text{ \& } \\ \eta_{\alpha_m} \upharpoonright (i+1) \in F_m(\eta_{\alpha_0} \upharpoonright (i+1), \dots, \eta_{\alpha_{m-1}} \upharpoonright (i+1), \eta_{\alpha_m} \upharpoonright i, \dots, \eta_{\alpha_n} \upharpoonright i) \}.$$

Note that if $i \in B_m$ then, as $\alpha_m^* \leq \alpha^* \leq \alpha_m$,

$$\eta_{\alpha_m} \upharpoonright (i+1) \in F_m(\eta_{\alpha_0} \upharpoonright (i+1), \dots, \eta_{\alpha_{m-1}} \upharpoonright (i+1), \eta_{\alpha_m} \upharpoonright i, \dots, \eta_{\alpha_n} \upharpoonright i) \subseteq \\ \subseteq F_m^*(\eta_{\alpha_0}, \dots, \eta_{\alpha_{m-1}}, \eta_{\alpha_m} \upharpoonright i) \in I_{\eta_{\alpha_m} \upharpoonright i}.$$

Hence we conclude that $B_m \subseteq B_m^*$ and therefore $B_m \in J$, what finishes the proof of the proposition. $\blacksquare$

Proposition 5.4 *Assume that $(T, \bar{\lambda}, \bar{I}) \in \mathcal{K}_{\mu, \delta}^{\text{id}}$, each ideal I_η (for $\eta \in T_i$, $i < \delta$) is $(|\delta| + |T_i|)^+$ -complete and $J \supseteq J_\delta^{\text{bd}}$ is an ideal on δ . Further suppose that a sequence $\bar{\eta} = \langle \eta_\alpha : \alpha < \lambda \rangle$ is super J -cofinal for $(T, \bar{\lambda}, \bar{I})$, λ is a regular cardinal greater than $|T|$ and a sequence $\langle \alpha_{\varepsilon, \ell} : \varepsilon < \lambda, \ell < n \rangle \subseteq \lambda$ is with no repetition and such that*

$$\alpha_{\varepsilon, 0} < \alpha_{\varepsilon, 1} < \dots < \alpha_{\varepsilon, n-1} \quad \text{for all } \varepsilon < \lambda.$$

Then for every $\varepsilon < \lambda$ large enough there is $a \in J$ such that

($\square$) if $i_\ell \in \delta \setminus a$ (for $\ell < n$), $i_0 \geq i_1 \geq \dots \geq i_{n-1}$ then

$$\begin{aligned} & (\exists^{I_{\eta_{\alpha_{\varepsilon, 0}} \restriction i_0}} \gamma_0) \dots (\exists^{I_{\eta_{\alpha_{\varepsilon, n-1}} \restriction i_{n-1}}} \gamma_{n-1}) \\ & (\exists^\lambda \zeta < \lambda) (\forall \ell < n) (\eta_{\alpha_{\zeta, \ell}} \restriction (i_\ell + 1) = \eta_{\alpha_{\varepsilon, \ell}} \restriction i_\ell \hat{\ } \langle \gamma_\ell \rangle). \end{aligned}$$

PROOF This is very similar to claim 4.3.1. First choose $\varepsilon_0 < \lambda$ such that for each $\varepsilon \in [\varepsilon_0, \lambda)$ and for every $i_0, \dots, i_{n-1} < \delta$ we have

$$(\exists^\lambda \zeta < \lambda) (\forall \ell < n) (\eta_{\alpha_{\zeta, \ell}} \restriction (i_\ell + 1) = \eta_{\alpha_{\varepsilon, \ell}} \restriction (i_\ell + 1))$$

(possible as $|T| < \text{cf}(\lambda) = \lambda$).

Now, for $\bar{i} = \langle i_\ell : \ell < n \rangle \subseteq \delta$ and $\bar{\nu} = \langle \nu_\ell : \ell < n \rangle$ such that $i_0 \geq i_1 \geq \dots \geq i_{n-1}$, $\nu_\ell \in T_{i_\ell}$ and $k < n$ we define a function $f_{\bar{i}, \bar{\nu}}^k : \prod_{\ell < k} \lambda_{\nu_\ell} \longrightarrow I_{\nu_k}$

(with a convention that $f_{\bar{i}, \bar{\nu}}^0$ is supposed to be a 0-place function, i.e. a constant) as follows

Let

$$B_{\bar{i}, \bar{\nu}} \stackrel{\text{def}}{=} \{ \langle \gamma_\ell : \ell < n \rangle \in \prod_{\ell < n} \lambda_{\nu_\ell} : (\exists^\lambda \zeta < \lambda) (\forall \ell < n) (\eta_{\alpha_{\zeta, \ell}} \restriction (i_\ell + 1) = \nu_\ell \hat{\ } \langle \gamma_\ell \rangle) \}.$$

If

$$(\spadesuit_{\bar{i}, \bar{\nu}}) \neg (\exists^{I_{\nu_0}} \gamma_0) \dots (\exists^{I_{\nu_{n-1}}} \gamma_{n-1}) (\langle \gamma_0, \dots, \gamma_{n-1} \rangle \in B_{\bar{i}, \bar{\nu}})$$

then $f_{\bar{i}, \bar{\nu}}^0, \dots, f_{\bar{i}, \bar{\nu}}^{n-1}$ are such that

$$\begin{aligned} (\diamond) \text{ if } \langle \gamma_0, \dots, \gamma_{n-1} \rangle \in B_{\bar{i}, \bar{\nu}} \\ \text{then } (\exists k < n) (\gamma_k \in f_{\bar{i}, \bar{\nu}}^k(\gamma_0, \dots, \gamma_{k-1})). \end{aligned}$$

Otherwise (i.e. if not $(\spadesuit_{\bar{i}, \bar{\nu}})$) the functions $f_{\bar{i}, \bar{\nu}}^k$ are constantly equal to $\emptyset$ (for $k < n$).

Next, for $k < n$, choose functions F_k such that if $\eta_0, \dots, \eta_k \in \lim_\delta(T)$, $i < \delta$ then

$$\begin{aligned} F_k(\eta_0, \dots, \eta_{k-1}, \eta_k \restriction i) = \\ \bigcup \{f_{\bar{i}, \bar{\nu}}^k(\eta_0(i_0), \dots, \eta_{k-1}(i_{k-1})) : \bar{i} = \langle i_\ell : \ell < n \rangle, \bar{\nu} = \langle \nu_\ell : \ell < n \rangle, \\ \delta > i_0 \geq \dots \geq i_k = i \geq i_{k+1} \geq \dots \geq i_{n-1} \\ \nu_\ell = \eta_\ell \restriction i_\ell \text{ for } \ell \leq k \text{ and} \\ \nu_\ell \in T_{i_\ell} \text{ for } k < \ell < n\}. \end{aligned}$$

Note that $F_k(\eta_0, \dots, \eta_{k-1}, \eta_k \restriction i)$ is a union of at most $|\delta| + |T_i|$ sets from the ideal $I_{\eta_k \restriction i}$ and hence $F_k(\eta_0, \dots, \eta_{k-1}, \eta_k \restriction i) \in I_{\eta_k \restriction i}$ (for each $\eta_0, \dots, \eta_k \in \lim_\delta(T)$, $i < \delta$). Thus, using the super J -cofinality of $\bar{\eta}$ we find $\alpha^* < \lambda$ such that

if $\alpha^* \leq \alpha_0 < \dots < \alpha_n < \lambda$
then the set

$$\{i < \delta : (\exists k < n)(\eta_{\alpha_k}(i) \in F_k(\eta_{\alpha_0}, \dots, \eta_{\alpha_{k-1}}, \eta_{\alpha_k}))\}$$

is in the ideal J .

Let $\varepsilon_1 > \varepsilon_0$ be such that for every $\varepsilon \in [\varepsilon_1, \lambda)$ we have $\alpha^* < \alpha_{\varepsilon,0} < \dots < \alpha_{\varepsilon,n-1}$.

Suppose now that $\varepsilon_1 < \varepsilon < \lambda$. By the choice of α^* we know that the set

$$a \stackrel{\text{def}}{=} \{i < \delta : (\exists \ell < n)(\eta_{\alpha_{\varepsilon,\ell}}(i) \in F_\ell(\eta_{\alpha_{\varepsilon,0}}, \dots, \eta_{\alpha_{\varepsilon,\ell-1}}, \eta_{\alpha_{\varepsilon,\ell}} \restriction i))\}$$

is in the ideal J . We are going to show that the assertion $(\square)$ holds for ε and a .

Suppose that $\bar{i} = \langle i_\ell : \ell < n \rangle \subseteq \delta \setminus a$, $i_0 \geq i_1 \geq \dots \geq i_{n-1}$. Let $\bar{\nu} = \langle \nu_\ell : \ell < n \rangle$, $\nu_\ell = \eta_{\alpha_{\varepsilon,\ell}} \restriction i_\ell$. If the condition $(\blacklozenge_{\bar{i}, \bar{\nu}})$ fails then we are done. So assume that it holds true. By the choice of the set a (and α^*) we have

$$(\forall \ell < n)(\eta_{\alpha_{\varepsilon,\ell}}(i_\ell) \notin F_\ell(\eta_{\alpha_{\varepsilon,0}}, \dots, \eta_{\alpha_{\varepsilon,\ell-1}}, \eta_{\alpha_{\varepsilon,\ell}} \restriction i_\ell))$$

what, by the definition of F_ℓ , implies that

$$(\forall \ell < n)(\eta_{\alpha_{\varepsilon,\ell}}(i_\ell) \notin f_{\bar{i}, \bar{\nu}}^\ell(\eta_{\alpha_{\varepsilon,0}}(i_0), \dots, \eta_{\alpha_{\varepsilon,\ell-1}}(i_{\ell-1}))).$$

By $(\diamond)$ we conclude that

$$\langle \eta_{\alpha_{\varepsilon,0}}(i_0), \dots, \eta_{\alpha_{\varepsilon,n-1}}(i_{n-1}) \rangle \notin B_{\bar{i}, \bar{\nu}}$$

and hence, by the definition of $B_{\bar{i}, \bar{\nu}}$,

$$\neg(\exists^\lambda \zeta)(\forall \ell < n)(\eta_{\alpha_{\zeta,\ell}} \restriction (i_\ell + 1) = \eta_{\alpha_{\varepsilon,\ell}} \restriction (i_\ell))$$

which contradicts the choice of ε_0 (remember $\varepsilon \geq \varepsilon_1 > \varepsilon_0$). ■

Definition 5.5 We say that a λ -marked Boolean algebra $(\mathbb{B}, \bar{y})$ has character n if

for every finite set $u \in [\lambda]^{<\omega}$ such that $\mathbb{B} \models \bigcap_{\alpha \in u} y_\alpha = \mathbf{0}$ there exist a subset $v \subseteq u$ of size $|v| \leq n$ such that $\mathbb{B} \models \bigcap_{\alpha \in v} y_\alpha = \mathbf{0}$.

Proposition 5.6 If a λ -marked Boolean algebra $(\mathbb{B}, \bar{y})$ is $(\theta, \text{not } \lambda)$ -Knaster (or other examples considered in the present paper) and $(\mathbb{B}, \bar{y})$ has character 2 then without loss of generality $(\mathbb{B}, \bar{y})$ is determined by a colouring on λ :

if $c : [\lambda]^2 \longrightarrow 2$ is such that

$$c(\{\alpha, \beta\}) = 0 \quad \text{iff} \quad \mathbb{B} \models y_\alpha \cap y_\beta = \mathbf{0}$$

then the algebra $\mathbb{B}$ is freely generated by $\{y_\alpha : \alpha < \lambda\}$ except that

if $c(\{\alpha, \beta\}) = 0$
then $y_\alpha \cap y_\beta = \mathbf{0}$. ■

Remark 5.7 These are nice examples.

Proposition 5.8 In all our results (like: 3.1 or 3.8), the marked Boolean algebra $(\mathbb{B}, \bar{y})$ which we get is actually of character 2 as long as any $(\mathbb{B}_\eta, \bar{y}_\eta)$ appearing in the assumptions (if any) is like that.

Then automatically the θ -Knaster property of the marked Boolean algebra $(\mathbb{B}, \bar{y})$ implies a stronger condition:

if $Z \in [\text{lg}(\bar{y})]^\theta$ then there is a set $Y \in [Z]^\theta$ such that $\{y_i : i \in Y\}$ generates a filter in $\mathbb{B}$. ■

Proposition 5.9 Let $(T, \bar{\lambda}, \bar{I}) \in \mathcal{K}_{\mu, \delta}^{\text{id}}$ be such that for each $\eta \in T$ the filter $(I_\eta)^c$ (dual to I_η) is an ultrafilter on $\text{succ}_T(\eta)$, and let J be an ideal on δ (extending J_δ^{bd}). If:

- (a) $\mathcal{C} = (T, \bar{\lambda}, \bar{\eta}, \langle (\mathbb{B}_\eta, \bar{y}_\eta) : \eta \in T \rangle)$ is a (δ, μ, λ) -constructor, the sequence $\bar{\eta}$ is stronger J -cofinal for $(T, \bar{\lambda}, \bar{I})$, $|T| < \text{cf}(\lambda) = \lambda$,
- (b) the sequence $\langle \alpha_{\varepsilon, \ell} : \varepsilon < \lambda, \ell < n \rangle \subseteq \lambda$ is with no repetition,
- (c) for each distinct $\eta, \nu \in T$ either the ideal I_η is $(2^{\lambda_\nu})^+$ -complete (which, of course, implies $\lambda_\eta > 2^{\lambda_\nu}$) or the ideal I_ν is $(2^{\lambda_\eta})^+$ -complete (it is enough if this holds true for η, ν such that $\text{lg}(\eta) = \text{lg}(\nu)$)

then for every large enough $\varepsilon < \lambda$ for J -almost all $i < \delta$ there are sets $X_\ell \in (I_{\eta_{\alpha_\varepsilon, \ell}} \restriction i)^+$ (for $\ell < n$) such that

$$(\forall \gamma_0 \in X_0) \dots (\forall \gamma_{n-1} \in X_{n-1}) (\exists^\lambda \zeta < \lambda) (\forall \ell < n) (\eta_{\alpha_\varepsilon, \ell} \restriction i \hat{=} \langle \gamma_\ell \rangle \triangleleft \eta_{\alpha_\zeta, \ell}).$$

Remark 5.9.A We can replace *stronger* by *big* and then omit being an ultrafilter.

PROOF First note that we may slightly re-enumerate the sequence $\langle \alpha_{\varepsilon, \ell} : \varepsilon < \lambda, \ell < n \rangle$ and we may assume that for each $\varepsilon < \lambda$

$$\alpha_{\varepsilon, 0} < \alpha_{\varepsilon, 1} < \dots < \alpha_{\varepsilon, n-1}.$$

Now, since $|T| < \text{cf}(\lambda) = \lambda$ we may apply claim 4.3.1 to

$$\langle \langle \alpha_{\varepsilon, \ell} : \ell < n \rangle : \varepsilon_0 \leq \varepsilon < \lambda \rangle$$

(we need to take ε_0 large enough to get the condition $(\hat{*})$ of the proof of 4.3). Consequently we may conclude that there is $\varepsilon_1 < \lambda$ such that for every $\varepsilon \in [\varepsilon_1, \lambda)$

$(\boxtimes_\varepsilon)$ for J -majority of $i < \delta$ we have

$$(\exists^{I_{\eta_{\alpha_\varepsilon, 0}} \restriction i} \gamma_0) \dots (\exists^{I_{\eta_{\alpha_\varepsilon, n-1}} \restriction i} \gamma_{n-1}) (\exists^\lambda \zeta < \lambda) (\forall \ell < n) (\eta_{\alpha_\zeta, \ell} \restriction (i+1) = \eta_{\alpha_\varepsilon, \ell} \restriction i \hat{=} \langle \gamma_\ell \rangle).$$

Now we would like to apply 1.2. We can not do this directly as we do not know if the cardinals $\lambda_{\eta_\varepsilon, \ell} \restriction i$ are decreasing (with ℓ). However the following claim helps us.

Claim 5.9.1 Suppose that $\lambda_0 < \lambda_1$ are cardinals and I_0, I_1 are maximal ideals on λ_0, λ_1 respectively. Assume that the ideal I_1 is $(\lambda_0)^+$ -complete and $\varphi(x, y)$ is a formula. Then

$$(\exists^{I_0} \gamma_0) (\exists^{I_1} \gamma_1) \varphi(\gamma_0, \gamma_1) \quad \Rightarrow \quad (\exists^{I_1} \gamma_1) (\exists^{I_0} \gamma_0) \varphi(\gamma_0, \gamma_1).$$

Why? First note that if I is a maximal ideal then the quantifiers $\exists^I$ and $\forall^I$ are equivalent. Suppose now that

$$(\exists^{I_0} \gamma_0) (\exists^{I_1} \gamma_1) \varphi(\gamma_0, \gamma_1).$$

This implies (as I_0, I_1 are maximal) that

$$(\forall^{I_0} \gamma_0) (\forall^{I_1} \gamma_1) \varphi(\gamma_0, \gamma_1).$$

Thus we have a set $a \in I_0$ and for each $\gamma \in \lambda_0 \setminus a$ we have a set $b_\gamma \in I_1$ such that

$$(\forall \gamma_0 \in \lambda_0 \setminus a)(\forall \gamma_1 \in \lambda_1 \setminus b_{\gamma_0})\varphi(\gamma_0, \gamma_1).$$

Let $b = \bigcup_{\gamma \in \lambda_0 \setminus a} b_\gamma$. As I_1 is $(\lambda_0)^+$ -complete the set b is in I_1 . Clearly

$$(\forall \gamma_1 \in \lambda_1 \setminus b)(\forall \gamma_0 \in \lambda \setminus a)\varphi(\gamma_0, \gamma_1)$$

which implies $(\exists^{I_1} \gamma_1)(\exists^{I_0} \gamma_0)\varphi(\gamma_0, \gamma_1)$, finishing the proof of the claim.

Now fix $\varepsilon > \varepsilon_1$ (ε_1 as chosen earlier). Take $i^* < \delta$ such that the elements of $\langle \eta_{\alpha_\varepsilon, \ell} \restriction i : \ell < n \rangle$ are pairwise distinct. Suppose that $i \in [i^*, \delta)$ is such that the formula of $(\boxtimes_\varepsilon)$ holds true. Let $\{k_\ell : \ell < n\}$ be an enumeration of n such that

$$\lambda_{\eta_{\alpha_\varepsilon, k_0} \restriction i} > \lambda_{\eta_{\alpha_\varepsilon, k_1} \restriction i} > \dots > \lambda_{\eta_{\alpha_\varepsilon, k_{n-1}} \restriction i}.$$

(Note that by the assumption (c) we know that all the $\lambda_{\eta_{\alpha_\varepsilon, k_\ell} \restriction i}$ are distinct, remember the choice of i^* .) Applying claim 5.9.1 we conclude that

$$(\exists^{I_{\eta_{\alpha_\varepsilon, k_0} \restriction i}} \gamma_{k_0}) \dots (\exists^{I_{\eta_{\alpha_\varepsilon, k_{n-1}} \restriction i}} \gamma_{k_{n-1}})(\exists^\lambda \zeta < \lambda)(\forall \ell < n)(\eta_{\alpha_\varepsilon, \ell} \restriction i \hat{=} \langle \gamma_\ell \rangle = \eta_{\alpha_\zeta, \ell} \restriction (i+1)).$$

But now we are able to use 1.2 and we get that there are sets $X_{k_\ell} \subseteq \lambda_{\eta_{\alpha_\varepsilon, k_\ell} \restriction i}$, $X_{k_\ell} \notin I_{\eta_{\alpha_\varepsilon, k_\ell} \restriction i}$ (for $\ell < n$) such that

$$\prod_{\ell < n} X_{k_\ell} \subseteq \{ \langle \gamma_0, \dots, \gamma_{n-1} \rangle : (\exists^\lambda \zeta < \lambda)(\forall \ell < n)(\eta_{\alpha_\varepsilon, \ell} \restriction i \hat{=} \langle \gamma_\ell \rangle = \eta_{\alpha_\zeta, \ell} \restriction (i+1)) \}$$

what is exactly what we need. ■

If we assume less completeness of the ideals I_η in 5.9 then still we may say something.

Proposition 5.10 *Let $\langle \sigma_i : i < \delta \rangle$ be a sequence of cardinals. Suppose that $T, \bar{\lambda}, \bar{I}, \bar{\eta}, J, \lambda, \mu, \delta$ and $\langle \alpha_{\varepsilon, \ell} : \varepsilon < \lambda, \ell < n \rangle$ are as in 5.9 but with condition (c) replaced by*

(c) $^-_{\langle \sigma_i : i < \delta \rangle}$ *if $\eta, \nu \in T_i$, $\eta \neq \nu$, $i < \delta$ then either the ideal I_η is $((\lambda_\nu)^{\sigma_i})^+ -$ complete or the ideal I_ν is $((\lambda_\eta)^{\sigma_i})^+ -$ complete.*

Then for every large enough $\varepsilon < \lambda$ for J -almost all $i < \delta$ there are sets $X_\ell \in [\lambda_{\eta_{\alpha_\varepsilon, \ell} \restriction i}]^{\sigma_i}$ (for $\ell < n$) such that

$$(\forall \gamma_0 \in X_0) \dots (\forall \gamma_{n-1} \in X_{n-1})(\exists^\lambda \zeta < \lambda)(\forall \ell < n)(\eta_{\alpha_\varepsilon, \ell} \restriction i \hat{=} \langle \gamma_\ell \rangle \triangleleft \eta_{\alpha_\zeta, \ell}).$$

PROOF The proof goes exactly as the one of 5.9, but instead of 1.2 we use 1.3. ■

Remark 5.11 1. Note that in the situation as in 5.9, we usually have that “ J -cofinal” implies “stronger J -cofinal” (see 2.7, 2.5).

2. The first assumption of 5.9 (ultrafilters) coupled with our normal completeness demands is a very heavy condition, but it has rewards.
3. A natural context here is when $\langle \mu_i : i \leq \kappa \rangle$ is a strictly increasing continues sequence of cardinals such that each μ_{i+1} is compact and $\mu = \mu_\kappa$. Then every μ_{i+1} -complete filter can be extended to an μ_{i+1} -complete ultrafilter. Moreover $2^\mu = \mu^+$ follows by Solovay [So74].

If for some function f from cardinals to cardinals, for each χ there is an algebra $\mathbb{B}_\chi$ of cardinality $f(\chi)$ which cannot be decomposed into $\leq \mu$ sets X_i each with some property $\mathbf{Pr}(\mathbb{B}_\chi, X_i)$ and if each μ_i is f -inaccessible

then we can find $T, \bar{I}, \bar{\lambda}$ as in 5.9 and such that $\eta \in T_i \Rightarrow \mu_i < \chi_\eta < \lambda_\eta < \mu_{i+1}$ and for $\eta \in T_i$ there is an algebra $\mathbb{B}_\eta$ with universe λ_η and the ideal I_η is χ_η -complete,

if $X \subseteq \mathbb{B}_\eta$ and $\mathbf{Pr}(\mathbb{B}_\eta, X)$ then $X \in I_\eta$

(compare 3.1) and $\lambda_\eta < \lambda_\nu \Rightarrow (2^{\lambda_\eta})^+ < \chi_\nu$. Now choosing cofinal $\bar{\eta}$ we may proceed as in earlier arguments.

4. It seems to be good for building nice examples, however we did not find the right question yet.
5. Central to our proofs is an assumption that

“ $\langle \alpha_{\zeta, \ell} : \zeta < \lambda, \ell < n \rangle \subseteq \lambda$ is a sequence with no repetition”,

i.e. we deal with λ disjoint n -tuples. This is natural as the examples constructed here are generated from $\{x_i : i < \lambda\}$ by finitary functions. One may ask what happens if we admit functions with, say, $\aleph_0$ places? We can still try to get for μ as above that:

($\boxtimes$) there is $h : [\mu^+]^2 \longrightarrow 2$ such that

if $\langle u_\varepsilon : \varepsilon < \lambda \rangle$ are pairwise disjoint, $u_\varepsilon = \{\alpha_{\varepsilon, \ell} : \ell < \ell^*\}$ is the increasing (with ℓ) enumeration, $\ell^* < \mu$ (ℓ^* infinite), for a sequence $\langle \nu_\ell : \ell < \ell^* \rangle \subseteq T_i$

$$B_{\langle \nu_\ell : \ell < \ell^* \rangle} \stackrel{\text{def}}{=} \{ \langle \eta_{\alpha_{\varepsilon, \ell}}(i) : \ell < \ell^* \rangle : (\exists \lambda \zeta < \lambda) (\forall \ell < \ell^*) (\eta_{\alpha_{\varepsilon, \ell}} \upharpoonright (i+1) = \eta_{\alpha_{\zeta, \ell}} \upharpoonright (i+1)) \},$$

for some $i^* < \delta$ there are no repetitions in $\langle \eta_{\alpha_{\varepsilon, \ell}} \restriction i^* : \ell < \ell^* \rangle$ and
 $h \restriction [u_\varepsilon]^2 \equiv 1$ (for each $\varepsilon < \lambda$)
 then there are $\alpha < \beta$ (really a large set of these) such that

$$h \restriction [u_\alpha \cup u_\beta]^2 \equiv 1.$$

The point is that we can deal with functions with infinitely many variables. Looking at previous proofs, “in stronger” we can get (for μ strong limit singular etc):

for α large enough
 for $i < \delta = \text{cf}(\mu)$ large enough

 we can defeat

$$(\dots (\forall^{I_{\eta_{\alpha_{\varepsilon, \ell}} \restriction i}} \gamma_\ell) \dots) (\langle \gamma_\ell : \ell < \ell^* \rangle \in B_{\langle \eta_{\alpha_{\varepsilon, \ell}} \restriction i : \ell < \ell^* \rangle})$$

but the duality of quantifiers fails, so the conclusion is only
 that

$$(\forall^J i < \delta) [\neg (\dots (\forall^{I_{\eta_{\alpha_{\varepsilon, \ell}} \restriction i}} \gamma_\ell) \dots)_{\ell < \ell^*} (\langle \eta_{\alpha_{\varepsilon, \ell}}(i) : \ell < \ell^* \rangle \notin B_{\langle \eta_{\alpha_{\varepsilon, \ell}} \restriction i : \ell < \ell^* \rangle})].$$

6. (no ultrafilters) If $I \supseteq J_\eta^{\text{bd}}$, δ is a regular cardinal, $\lambda_\eta = \lambda_{\text{lg}(\eta)}$ and for each $u \in [T_i]^{<|\delta|\chi}$, $i < \delta$ the free product $\bigstar_{\eta \in u} \mathbb{B}_\eta$ satisfies the λ -cc then we can show that the algebra $\mathbb{B}_{<\chi}^{\text{red}}$ satisfies the λ -cc too,

where for a cardinal χ the algebra $\mathbb{B}_{<\chi}^{\text{red}}$ is the Boolean algebra freely generated by

$$\begin{aligned} \{ \bigcap_{\alpha \in u} x_\alpha^{t(\alpha)} : t : u \longrightarrow 2, u \in [\lambda]^{<\delta}, h \restriction [u \cap t^{-1}[1]]^2 \equiv 1 \text{ and} \\ |u| < \chi \text{ and} \\ (\exists i < \delta) (\text{the mapping } \alpha \mapsto \eta_\alpha(i) \text{ is one-to-one (for } \alpha \in u)) \\ (\exists i < \delta) (\exists \alpha \in u) (\forall j \in (i, \delta)) (\forall \beta \in u) (f_\alpha(j) \leq f_\beta(j)) \}. \end{aligned}$$

[Note that if $\chi \leq \text{cf}(\delta)$ simpler.]

* * * * *

Now we will deal with an additional demand that the algebra $\mathbb{B}^{\text{red}}$ satisfies $|\delta|^+$ -cc (or even has the $|\delta|^+$ -Knaster property). Note that the demand of $|\delta|$ -cc does not seem to be reasonable: if every $\bar{y}_\eta$ has two disjoint members (and every node $t \in T$ is an initial segment of a branch through T) then we can find δ branches which, if in $\{\eta_\alpha : \alpha < \lambda\}$, give δ pairwise disjoint elements. *Moreover:*

for each $\nu \in T_\ell$ let $A_\nu = \{\eta_\alpha(i) : \eta_\alpha \restriction i = \nu\}$ and

$$a_\alpha = \{i < \delta : (\exists \beta \in A_{\eta_\alpha \restriction i})(\mathbb{B}_{\eta_\alpha \restriction i} \models y_{\eta_\alpha(i)} \cap y_\beta = \mathbf{0})\}.$$

So if $\mathbb{B}^{\text{red}} \models \sigma\text{-cc}$ then $(\forall \alpha < \lambda)(|a_\alpha| < \sigma)$.

Definition 5.12 Let $(T, \bar{\lambda}) \in \mathcal{K}_{\mu, \delta}$ and let $\bar{\eta} = \langle \eta_\alpha : \alpha < \lambda \rangle \subseteq \lim_\delta(T)$. We say that $\bar{\eta}$ is hereditary θ -free if for every $Y \in [\lambda]^\theta$ there are $Z \in [Y]^\theta$ and $i < \delta$ such that

$$(\forall \alpha, \beta \in Z)(\alpha \neq \beta \Rightarrow [\eta_\alpha \restriction i = \eta_\beta \restriction i \ \& \ \eta_\alpha(i) \neq \eta_\beta(i)]).$$

Proposition 5.13 Assume that $\mathcal{C} = (T, \bar{\lambda}, \bar{\eta}, \langle (\mathbb{B}_\eta, \bar{y}_\eta) : \eta \in T \rangle)$ is a (δ, μ, λ) -constructor. If $\bar{\eta}$ is hereditary θ -free, each algebra $\mathbb{B}_\eta$ has the θ -Knaster property and θ is regular then the algebra $\mathbb{B}^{\text{red}}(\mathcal{C})$ has the θ -Knaster property.

PROOF The same as for 3.8. Note that the proof there shows actually that,

if $(\forall \alpha < \theta)(|\alpha|^{\delta} < \theta = \text{cf}(\theta))$
then $\bar{\eta}$ is θ -hereditary free. ■

Proposition 5.14 Assume that $(T, \bar{\lambda}) \in \mathcal{K}_{\mu, \delta}$, $\bar{\eta} = \langle \eta_\alpha : \alpha < \lambda \rangle \subseteq \lim_\delta(T)$, λ is a regular cardinal. Further suppose that

- (a) $(\forall \alpha < \theta)(|\alpha|^{<\delta} < \theta = \text{cf}(\theta))$, $\delta < \theta$, J is an ideal on δ extending J_δ^{bd} and
- (b) the sequence $\bar{\eta}$ is $<_J$ -increasing and one of the following conditions is satisfied:
 - (α) $\bar{\eta}$ is $<_J$ -cofinal in $\prod_{i < \delta} \lambda_i / J$, λ_i are regular cardinals above θ (at least for J -majority of $i < \delta$), $\{\alpha < \lambda : \text{cf}(\alpha) = \theta\} \in I[\lambda]$ and $\lambda_\eta = \lambda_{\text{lg}(\eta)}$;
 - (β) there are a sequence $\langle C_\alpha : \alpha < \lambda \rangle$ of subsets of λ , a closed unbounded subset E of λ and $i^* < \delta$ such that
 1. $C_\alpha \subseteq \alpha$, $\text{otp}(C_\alpha) \leq \theta$,
 2. if $\beta \in C_\alpha$ then $C_\beta = C_\alpha \cap \beta$ and $\eta_\beta \restriction [i^*, \delta) < \eta_\alpha \restriction [i^*, \delta)$,
 3. if $\alpha \in E$ and $\text{cf}(\alpha) = \theta$ then $\alpha = \sup(C_\alpha)$.

Then there is $A \in [\lambda]^\lambda$ such that the restriction $\bar{\eta} \upharpoonright A$ is θ -hereditary free.

PROOF First let us assume that the case (β) of the clause **(b)** of the assumptions holds.

Claim 5.14.1 Suppose that $Y \in [E]^\theta$. Then

1. $(\exists Z \in [Y]^\theta)(\exists i^\otimes)(\text{the sequence } \langle f_{\beta_\varepsilon}(i^\otimes) : \varepsilon \in Z \rangle \text{ is strictly increasing}).$
2. If additionally $J = J_\delta^{\text{bd}}$ then

$(\exists Z \in [Y]^\theta)(\exists i^\otimes < \delta)(\text{the sequence } \langle \eta_\beta \upharpoonright [i^\otimes, \delta) : \beta \in Z \rangle \text{ is strictly increasing}).$

Why? Suppose $Y \in [E]^\theta$. Without loss of generality we may assume that $\text{otp}(Y) = \theta$. Let $\alpha = \sup(Y)$. So $\alpha \in E$, $\text{cf}(\alpha) = \theta$ and hence C_α is unbounded in α . Let $C_\alpha = \langle \alpha_\varepsilon : \varepsilon < \theta \rangle$ be the increasing enumeration. Clearly the set

$$A \stackrel{\text{def}}{=} \{\varepsilon < \theta : [\alpha_\varepsilon, \alpha_{\varepsilon+1}) \cap Y \neq \emptyset\}$$

is unbounded in θ . For $\varepsilon \in A$ choose $\beta_\varepsilon \in [\alpha_\varepsilon, \alpha_{\varepsilon+1}) \cap Y$. Then

$$(\exists a_\varepsilon \in J)(\eta_{\alpha_\varepsilon} \upharpoonright (\delta \setminus a_\varepsilon) \leq \eta_{\beta_\varepsilon} \upharpoonright (\delta \setminus a_\varepsilon) < \eta_{\alpha_{\varepsilon+1}} \upharpoonright (\delta \setminus a_\varepsilon)).$$

Now choose $i_\varepsilon \in \delta \setminus a_\varepsilon$, $i_\varepsilon > i^*$ and find $B \in [A]^\theta$ such that

$$\varepsilon \in B \quad \Rightarrow \quad i_\varepsilon = i^\otimes.$$

Easily, by the assumption $(\beta)(2)$, this $i^\otimes$ and $Z = \{\beta_\varepsilon : \varepsilon \in B\}$ are as required in 5.14.1(1).

If additionally we know that $J = J_\delta^{\text{bd}}$ then for some $B \in [A]^\theta$ we have

$$(\exists i^\otimes \in [i^*, \delta))(\varepsilon \in B \quad \Rightarrow \quad a_\varepsilon \subseteq i^\otimes)$$

and hence the sequence $\langle f_{\beta_\varepsilon} \upharpoonright [i^\otimes, \delta) : \varepsilon \in B \rangle$ is as required in 5.14.1(2) (remember $(\beta)(2)$).

But now, using $i^\otimes$ given by 5.14.1 we may deal with the sequence $\langle f_{\beta_\varepsilon} \upharpoonright (i^\otimes + 1) : \varepsilon \in B \rangle$ and using the old proof (see 3.8) on the tree $\bigcup_{i \leq i^\otimes} T_i$ (note that we may apply the assumption **(a)** to arguments like there) we may get the desired conclusion. This finishes the case when (β) of **(b)** holds true.

Now, assume that the case (α) of the clause **(b)** of the assumptions holds. We reduce this case to the previous one (using cofinality).

Take $\bar{C}$, E witnessing that the set $\{\alpha < \lambda : \text{cf}(\alpha) = \theta\}$ is in $I[\lambda]$ and build a $<_J$ -increasing sequence $\bar{\eta}' = \langle \eta'_\alpha : \alpha < \lambda \rangle \subseteq \prod_{i < \delta} \lambda_i$ such that $\eta'_\alpha > \eta_\alpha$ and $\bar{\eta}'$ satisfies the clause (β) of **(b)** for $\bar{C}$, E . [The construction of η'_α is by induction on $\alpha < \lambda$. Suppose that we have defined η'_β for $\beta < \alpha$. Now, at the stage α of the construction, we first choose $\eta_\alpha^0 \in \prod_{i < \delta} \lambda_i$ such that

$$(\forall \beta < \alpha)(\eta'_\beta <_J \eta_\alpha^0).$$

This is possible since the condition (α) implies that $\lambda = \text{tcf}(\prod_{i < \delta} \lambda_i / J)$, $\alpha < \lambda$. Now we put for $i < \delta$:

$$\eta'_\alpha(i) = \max \{ \eta_\alpha^0(i), \eta_\alpha(i) + 1, \sup \{ \eta'_\gamma(i) + 1 : \gamma \in C_\alpha \} \}.$$

Now one can check that this $\bar{\eta}'$ is as required.]

Now we use the fact that $\bar{\eta}$ is cofinal. The set

$$E' = \{ \gamma \in E : (\forall \alpha < \gamma)(\exists \beta < \gamma)(\eta'_\alpha <_J \eta_\beta) \}$$

is a club of λ . Look at $\bar{\eta} \restriction E'$. Suppose that $Y \in [E']^\theta$. Without loss of generality we may assume that $\text{otp}(Y) = \theta$ and let $\alpha = \sup(Y)$. By induction on $\varepsilon < \theta$ choose $\alpha_\varepsilon < \beta_\varepsilon < \gamma_\varepsilon$ such that

$$\begin{aligned} & \beta_\varepsilon \in Y, \alpha_\varepsilon \in C_\alpha, \gamma_\varepsilon \in C_\alpha, \eta'_{\alpha_\varepsilon} <_J \eta_{\beta_\varepsilon} <_J \eta'_{\gamma_\varepsilon} \text{ and} \\ & \text{if } \zeta < \varepsilon \text{ then } \gamma_\zeta < \alpha_\varepsilon. \end{aligned}$$

Next choose $i_\varepsilon > i^*$ such that

$$\eta'_{\alpha_\varepsilon}(i_\varepsilon) < \eta_{\beta_\varepsilon}(i_\varepsilon) < \eta'_{\gamma_\varepsilon}(i_\varepsilon).$$

We may assume that $i_\varepsilon = i^\otimes$ for all $\varepsilon < \theta$. Now, as $\bar{\eta}'$ obeys $\bar{C}$, we have

$$\zeta < \varepsilon \Rightarrow \eta'_{\gamma_\zeta}(i^\otimes) < \eta'_{\alpha_\varepsilon}(i^\otimes)$$

and hence we conclude that the sequence $\langle \eta_{\beta_\varepsilon}(i^\otimes) : \varepsilon < \theta \rangle$ is strictly increasing. Now we may finish the proof like earlier. ■

Conclusion 5.15 *If μ is a strong limit singular cardinal, $2^\mu = \mu^+ = \lambda$ and $\neg(\exists 0^\#)$ or at least*

$$\{ \delta < \mu^+ : \text{cf}(\delta) = (2^{<\text{cf}(\mu)})^+ \} \in I[\lambda]$$

then there is a $(\text{cf}(\mu), \mu, \lambda)$ -constructor $\mathcal{C}$ such that the algebra $\mathbb{B}^{\text{red}}(\mathcal{C})$ has the $(2^{<\text{cf}(\mu)})^+$ -Knaster property, its counterpart $\mathbb{B}^{\text{green}}(\mathcal{C})$ is λ -cc and the free product is not λ -cc.

[Note that if GCH holds then $(2^{<\text{cf}(\mu)})^+ = (\text{cf}(\mu))^+$ so the problem is closed then.]

PROOF Like 4.4 using 5.14, 5.13 instead of 2.8, 3.8. ■

6 The use of pcf

Assuming that $2^{<\kappa}$ is much larger than $\kappa = \text{cf}(\kappa)$ ($= \text{cf}(\mu) < \mu$) we may still want to have examples with the $(\kappa^+, \text{not } \lambda)$ -Knaster property and the non-multiplicativity. Here 5.15 does not help if GCH holds on an end segment of the cardinals (and $\neg(\exists 0^\#)$). We try to remedy this.

It is done inductively. So 6.3 uses $\text{cf}(\mu) = \aleph_0$ just to start the induction. We can phrase (a part of) it without this assumption but in applications we use it for $\text{cf}(\mu) = \aleph_0$. Also 6.3(b) really needs this condition (otherwise we would have to assume that $(\forall \alpha < \theta)(|\alpha|^{<\delta} < \mu)$). This result says that, if $\text{cf}(\mu) = \aleph_0$, then we have gotten the θ -Knaster property for *every* regular cardinal $\theta \in \mu \setminus \kappa^+$.

Definition 6.1 1. By $\mathcal{K}_{\text{wmk}}$ we will denote the class of all tuples $(\theta, \lambda, \chi, J)$ such that $\theta < \lambda$, χ are regular cardinals, J is a χ -complete ideal on λ and there is a (λ, χ) -well marked Boolean algebra $(\mathbb{B}, \bar{y}, J)$ (see 3.2) such that the algebra $\mathbb{B}$ satisfies the θ -Knaster property (wmk stays for “**well marked Knaster**”).

When we write $(\theta, \lambda) \in \mathcal{K}_{\text{wmk}}$ we really mean $(\theta, \lambda, \lambda, J_\lambda^{\text{bd}}) \in \mathcal{K}_{\text{wmk}}$ (what means just that there exists a (θ, λ) -Knaster marked Boolean algebra).

2. By $\mathcal{K}_{\text{smk}}$ (smk is for “**sequence marked Knaster**”) we will denote the class of all triples (θ, λ, χ) of cardinals such that $\theta < \lambda$ are regular and there is a sequence $\langle (\mathbb{B}_\alpha, \bar{y}^\alpha) : \alpha < \chi \rangle$ of λ -marked Boolean algebras such that (for $\alpha < \chi$) the algebras $\mathbb{B}_\alpha$ have the θ -Knaster property, $\bar{y}^\alpha = \langle y_i^\alpha : i < \lambda \rangle$ and

if $n < \omega$, $\alpha_0 < \dots < \alpha_{n-1} < \chi$ and $\beta_{\varepsilon, \ell} < \lambda$ for $\varepsilon < \lambda$, $\ell < n$ are such that $(\forall \varepsilon_1 < \varepsilon_2 < \lambda)(\forall \ell < n)(\beta_{\varepsilon_1, \ell} < \beta_{\varepsilon_2, \ell})$ then there are $\varepsilon_1 < \varepsilon_2 < \lambda$ such that

$$\ell < n \quad \Rightarrow \quad \mathbb{B}_{\alpha_\ell} \models “y_{\beta_{\varepsilon_1, \ell}}^{\alpha_\ell} \cap y_{\beta_{\varepsilon_2, \ell}}^{\alpha_\ell} = \mathbf{0}”.$$

Remark 6.2 1. On some closure properties of $\mathcal{K}_{\text{wmk}}^\theta \stackrel{\text{def}}{=} \{\lambda : (\theta, \lambda) \in \mathcal{K}_{\text{wmk}}\}$ under pcf see 3.12: if $\lambda_i \in \mathcal{K}_{\text{wmk}}^\theta$ (for $i < \delta$), $\lambda_i > \max \text{pcf}\{\lambda_j : j < i\}$ and $\lambda \in \text{pcf}\{\lambda_i : i < \delta\}$ and $(\forall \alpha < \theta)(|\alpha|^{|\delta|} < \theta)$ then $\lambda \in \mathcal{K}_{\text{wmk}}^\theta$.

2. We can replace θ by a set Θ of such cardinals, no real difference. And thus we may consider the class $\mathcal{K}_{\text{wmk}}^*$ of all tuples $(\Theta, \lambda, \chi, J)$ such that there exists a (λ, χ) -well marked Boolean algebra $(\mathbb{B}, \bar{y}, J)$ with

$$(\forall \theta \in \Theta)(\mathbb{B} \text{ satisfies the } \theta\text{-Knaster property}).$$

Proposition 6.3 *Assume that μ is a strong limit singular cardinal, $\aleph_0 = \text{cf}(\mu) < \mu$ and $\lambda = 2^\mu = \mu^+$.*

- (a) *If $(\forall \alpha < \theta)(|\alpha|^{\text{cf}(\mu)} < \theta = \text{cf}(\theta) < \lambda)$
then $(\theta, \lambda) \in \mathcal{K}_{\text{wmk}}$. Moreover $(\theta, \lambda, 2^\lambda) \in \mathcal{K}_{\text{smk}}$.*
- (b) *If $\text{cf}(\mu) < \theta = \text{cf}(\theta) < \mu$ and $\{\alpha < \lambda : \text{cf}(\alpha) = \theta\} \in I[\lambda]$
then $(\theta, \lambda) \in \mathcal{K}_{\text{wmk}}$. Moreover $(\theta, \lambda, 2^\lambda) \in \mathcal{K}_{\text{smk}}$.*

PROOF This is similar to previous proofs and the first parts of 6.3(a), (b) follow from what we have done already: 6.3(a) is an obvious modification of 3.11; 6.3 is similar, but based on 5.13, 5.14 (and 2.8, 3.7) (see below). What we actually have to prove are the “moreover” parts. We will sketch the proof of it for clause (b) only, modifying the proof of 4.4.

As in 4.4 we choose a function $h : \text{cf}(\mu) \longrightarrow \omega$ such that for each $n \in \omega$ the preimage $h^{-1}[\{n\}]$ is unbounded (in $\text{cf}(\mu)$). Next we take an increasing equence $\langle \mu_i : i < \text{cf}(\mu) \rangle$ of regular cardinals such that $\mu = \sum_{i < \delta} \mu_i$. Finally (like in 4.4) we construct $\lambda_i, \chi_i, (\mathbb{B}_i, \bar{y}_i)$ and I_i such that for $i < \text{cf}(\mu)$:

1. $\lambda_i, \chi_i < \mu$ are regular cardinals,
2. $\lambda_i > \chi_i \geq \prod_{j < i} \lambda_j + \mu_i$, $\chi_0 > \theta + \mu_0$,
3. I_i is a χ_i^+ -complete ideal on λ_i ,
4. $(\mathbb{B}_i, \bar{y}_i)$ is a λ_i -marked Boolean algebra such that
if $n = h(i)$ and the set $Y \subseteq (\lambda_i)^{n+1}$ is such that

$$(\exists^{I_i} \gamma_0) \dots (\exists^{I_i} \gamma_n) (\langle \gamma_0, \dots, \gamma_n \rangle \in Y)$$

then for some $\gamma'_\ell, \gamma''_\ell < \lambda_i$ (for $\ell \leq n$) we have

$$\langle \gamma'_\ell : \ell \leq n \rangle, \langle \gamma''_\ell : \ell \leq n \rangle \in Y \quad \text{and for all } \ell \leq n$$

$$\mathbb{B}_i \models y_{\gamma'_\ell}^i \cap y_{\gamma''_\ell}^i = \mathbf{0},$$

5. each algebra $\mathbb{B}_i$ satisfies the θ -Knaster condition,
6. for $\xi < \lambda_i$ the set $[\xi, \lambda_i)$ is not in the ideal I_i .

Note that the last requirement is new here. Though we cannot demand that the ideals I_i extend $I_{\lambda_i}^{\text{bd}}$, the condition (6) above is satisfied in our standard construction. Note that the ideal from 3.1 has this property if λ there is regular. Moreover it is preserved when the (finite) products of ideals (as in 4.2) are considered. Also, if I is an ideal on λ , $A_0 \in I$ is such that $|\lambda \setminus A_0|$ is minimal and $A_1 \in I^+$ is such that $|A_1|$ is minimal then we can use either $I \restriction A_0$ or $I \restriction A_1$. All relevant information is preserved then (in the first case the condition (6) holds in the second $J_\lambda^{\text{bd}} \subseteq I$ – under suitable renaming). Now we put $T = \bigcup_{i < \delta} \prod_{j < i} \lambda_j$, $\mathbb{B}_\eta = \mathbb{B}_{\text{lg}(\eta)}$, $\bar{y}_\eta = \bar{y}_{\text{lg}(\eta)}$, $I_\eta = I_{\text{lg}(\eta)}$. Applying

2.8 we find a stronger J_δ^{bd} -cofinal sequence $\bar{\eta} = \langle \eta_\alpha : \alpha < \lambda \rangle$ for $(T, \bar{\lambda}, \bar{I})$. Due to the requirement (6) above we may additionally demand that $\bar{\eta}$ is $<_{J_{\text{cf}(\mu)}^{\text{bd}}}$ -increasing cofinal in $\prod_{i < \text{cf}(\mu)} \lambda_i / J_{\text{cf}(\mu)}^{\text{bd}}$. Let $\langle B_\xi : \xi < 2^\lambda \rangle$ be a sequence

of pairwise almost disjoint elements of $[\lambda]^\lambda$ (i.e. $|B_\xi \cap B_\zeta| < \lambda$ for distinct $\xi, \zeta < 2^\lambda$). For each $\xi < 2^\lambda$ we may apply 5.14 (the version of **(b)**(α)) to the sequence $\langle \eta_\alpha : \alpha \in B_\xi \rangle$ and we find $A_\xi \in [B_\xi]^\lambda$ such that each sequence $\langle \eta_\alpha : \alpha \in A_\xi \rangle$ is θ -hereditary free. Let

$$\mathbb{B}_\xi^* = \mathbb{B}^{\text{red}}(T, \bar{\lambda}, \langle \eta_\alpha : \alpha \in A_\xi \rangle, \langle (\mathbb{B}_\eta, \bar{y}_\eta) : \eta \in T \rangle), \quad \bar{x}_\xi = \langle x_\alpha^{\text{red}} : \alpha \in A_\xi \rangle.$$

Of course each $\mathbb{B}_\xi^*$ is a subalgebra of $\mathbb{B}^{\text{red}}(T, \bar{\lambda}, \bar{\eta}, \langle (\mathbb{B}_\eta, \bar{y}_\eta) : \eta \in T \rangle)$ (generated by $\bar{x}_\xi$). By 5.13 and 3.7 we know that the marked Boolean algebras $(\mathbb{B}_\xi^*, \bar{x}_\xi)$ are $(\theta, \text{not}\lambda)$ -Knaster. To show that they witness $(\theta, \lambda, 2^\lambda) \in \mathcal{K}_{\text{smk}}$ suppose that $n < \omega$, $\xi_0, \dots, \xi_{n-1} < 2^\lambda$, $\beta_{\varepsilon, \ell} < \lambda$ (for $\varepsilon < \lambda$, $\ell < n$) are such that

$$(\forall \varepsilon_1 < \varepsilon_2 < \lambda)(\forall \ell < n)(\beta_{\varepsilon_1, \ell} < \beta_{\varepsilon_2, \ell})$$

and of course $\{\beta_{\varepsilon, \ell} : \varepsilon < \lambda\} \subseteq A_{\xi_\ell}$. Since A_{ξ_ℓ} are almost disjoint we may assume that

$$(\forall \varepsilon_1, \varepsilon_2 < \lambda)(\forall \ell_1 < \ell_2 < n)(\beta_{\varepsilon_1, \ell_1} \neq \beta_{\varepsilon_2, \ell_2}).$$

Further we may assume that we have $i^* < \text{cf}(\mu)$ such that for each $\varepsilon < \lambda$ the sequences $\eta_{\beta_{\varepsilon, \ell}} \restriction i^*$ for $\ell < n$ are pairwise distinct.

By the choice of $\bar{\eta}$, T , $\bar{\lambda}$ etc we may apply 4.3.1 and conclude that for all sufficiently large $\varepsilon < \lambda$ the set

$$Z_\varepsilon = \{i < \text{cf}(\mu) : \neg(\exists^{I_{\eta_{\beta_{\varepsilon, 0}}}} \gamma_0) \dots (\exists^{I_{\eta_{\beta_{\varepsilon, n-1}}}} \gamma_{n-1})(\exists^\lambda \zeta)(\forall \ell < n) \\ (\eta_{\beta_{\varepsilon, \ell}} \restriction (i+1) = (\eta_{\beta_{\varepsilon, \ell}} \restriction i)^\wedge \langle \gamma_\ell \rangle)\}$$

is in the ideal $J_{\text{cf}(\mu)}^{\text{bd}}$. Take one such ε . Choosing $i \in \text{cf}(\mu) \setminus Z_\varepsilon$, $i > i^*$ such that $h(i) = n$ we may follow exactly as in the last part of the proof of 4.3 and we find $\varepsilon_0, \varepsilon_1 < \lambda$ such that for each $\ell < n$

$$\eta_{\beta_{\varepsilon_0, \ell}} \restriction i = \eta_{\beta_{\varepsilon_1, \ell}} \restriction i \text{ but} \\ \mathbb{B}_{\eta_{\beta_{\varepsilon_0, \ell}}} \restriction i \models y_{\eta_{\beta_{\varepsilon_0, \ell}} \restriction (i+1)} \cap y_{\eta_{\beta_{\varepsilon_1, \ell}} \restriction (i+1)} = \mathbf{0}$$

which implies that

$$(\forall \ell < n)(\mathbb{B}_{\xi_\ell}^* \models x_{\beta_{\varepsilon_0, \ell}}^{\text{red}} \cap x_{\beta_{\varepsilon_1, \ell}}^{\text{red}} = \mathbf{0}).$$

■

Proposition 6.4 *Assume that*

- (a) $\langle \lambda_i : i < \delta \rangle$ *is an increasing sequence of regular cardinals such that*
 $\delta < \lambda_0$, $\lambda_i > \max \text{pcf}\{\lambda_j : j < i\}$ *(the last is our natural assumption);*
- (b) $\aleph_0 < \theta = \text{cf}(\theta) < \bigcup_{i < \delta} \lambda_i$ *(naturally we assume just* $\text{cf}(\theta) = \theta < \lambda_0$ *)*
- (c) $\lambda = \max \text{pcf}\{\lambda_i : i < \delta\}$
- (d) $(\theta, \lambda_i, \max \text{pcf}\{\lambda_j : j < i\}) \in \mathcal{K}_{\text{smk}}$
- (e) *for each* $\tau \in \{\lambda\} \cup \bigcup_{\alpha < \delta} \text{pcf}\{\lambda_i : i < \alpha\}$ *we have*

$$\{\xi < \tau : \text{cf}(\xi) = \theta\} \in I[\tau]$$

or at least

for some $\bar{f}^\tau = \langle f_\varepsilon^\tau : \varepsilon < \tau \rangle$, $<_{J=\tau}$ -*increasing cofinal in*
 $\prod_{i < \alpha} \lambda_i / J_{=\tau}$ *we have:*

$$\gamma < \tau \ \& \ \text{cf}(\gamma) = \theta \quad \Rightarrow \quad f_\gamma^\tau \text{ is good in } \bar{f}^\tau$$

(see [Sh 345a], [Sh 355], section 1 (see 1.6(1) and then [MgSh 204]),

- (f) $|\text{pcf}\{\lambda_i : i < \delta\}| < \theta$ *or at least for each* $\alpha < \delta$ *we have* $|\text{pcf}\{\lambda_i : i < \alpha\}| < \theta$.

Then $(\theta, \lambda) \in \mathcal{K}_{\text{wmk}}$. Moreover $(\theta, \lambda, \chi) \in \mathcal{K}_{\text{smk}}$ *provided there is an almost disjoint family of size* χ *in* $[\lambda]^\lambda$. *We may get algebras* $\mathbb{B}^{\text{red}}$, $\mathbb{B}^{\text{green}}$ *as in main constructions such that*

$$\mathbb{B}^{\text{red}} \models \theta\text{-Knaster}, \quad \mathbb{B}^{\text{green}} \models \lambda\text{-cc} \quad \text{and} \quad \mathbb{B}^{\text{red}} * \mathbb{B}^{\text{green}} \models \neg \lambda\text{-cc}.$$

Remark 6.4.A: Continues also the proof of 3.5 of [Sh:g].

PROOF The main difficulty of the proof will be to construct a hereditary θ -free $<_{J_{<\lambda}}$ -increasing sequence $\bar{\eta} = \langle \eta_\alpha : \alpha < \lambda \rangle \subseteq \prod_{i < \delta} \lambda_i$. This is done in the claim below. For the notation used there let us note that if $\alpha \leq \delta$ is a limit ordinal, $\tau \in \text{pcf}\{\lambda_i : i < \alpha\}$ then $J_{=\tau}[\{\lambda_i : i < \alpha\}] = J_\tau^\alpha$ is the ideal on α generated by

$$J_{<\tau}[\{\lambda_i : i < \alpha\}] \cup \{\alpha \setminus \text{b}_\tau[\{\lambda_i : i < \alpha\}]\}.$$

So in particular $\text{tcf}(\prod_{i < \alpha} \lambda_i / J_\tau^\alpha) = \tau$.

Claim 6.4.1 *There exists a tree $T \subseteq \bigcup_{i < \delta} \prod_{j < i} \lambda_j$ such that $\lim_\delta(T)$ is θ -hereditary free (and $<_{J_{<\lambda}}$ -cofinal).*

Moreover for each $\alpha < \delta$ the size of T_α is $\leq \max \text{pcf}\{\lambda_i : i < \alpha\}$.

Why? For a limit ordinal $\alpha \leq \delta$ and $\tau \in \text{pcf}\{\lambda_i : i \leq \alpha\}$ (if $\alpha = \delta$ then $\tau = \lambda$) choose a $<_{J_\tau^\alpha}$ -increasing sequence $\bar{f}^{\alpha, \tau} = \langle f_\zeta^{\alpha, \tau} : \zeta < \tau \rangle \subseteq \prod_{i < \alpha} \lambda_i$ cofinal in $\prod_{i < \alpha} \lambda_i / J_\tau^\alpha$ and such that

($\tilde{\otimes}$) if $\zeta < \tau$, $\text{cf}(\zeta) = \theta$

then for some unbounded set $Y_\zeta \subseteq \zeta$ (for simplicity consisting of successor ordinals) and a sequence $\bar{s}^\tau = \langle s_\xi^\tau : \xi \in Y_\zeta \rangle \subseteq J_\tau^\alpha$ we have

$$[\xi_1, \xi_2 \in Y_\zeta \ \& \ \xi_1 < \xi_2 \ \& \ i \in \alpha \setminus (s_{\xi_1}^\tau \cup s_{\xi_2}^\tau)] \Rightarrow f_{\xi_1}^{\alpha, \tau}(i) < f_{\xi_2}^{\alpha, \tau}(i).$$

[Why we can demand ($\tilde{\otimes}$)? If in the assumption (e) the first part is satisfied then we follow similarly to the proof of 5.14, compare [Sh 355], 1.5A, 1.6, pp 51–52. If we are in the case of “at least” then this is exactly the meaning of goodness.] Further we may demand that the sequence $\bar{f}^{\alpha, \tau}$ is b continuous:

($\tilde{\oplus}$) if $|\delta| < \text{cf}(\zeta) < \lambda_0$, $\zeta < \tau$

then $f_\zeta^{\alpha, \tau}(i) = \min\{\bigcup_{\xi \in C} f_\xi^{\alpha, \tau}(i) : C \text{ is a club of } \zeta\}$

[compare the proof of 3.4 of [Sh 345a], pp 25–26].

For a limit ordinal $\alpha < \delta$ we define

$$T_\alpha^0 = \{f \in \prod_{i < \alpha} \lambda_i : \text{(a) } f = \max\{f_{\zeta_\ell}^{\alpha, \tau_\ell} : \ell < n\} \text{ for some}$$

$$n < \omega, \tau_\ell \in \text{pcf}\{\lambda_i : i < \alpha\}, \zeta_\ell < \tau_\ell$$

(b) for every $\tau \in \text{pcf}\{\lambda_i : i < \alpha\}$ if $\tau = \lambda$ or $\alpha < \delta$ then there is $\zeta_f(\tau) < \tau$ such that

$$f_{\zeta_f(\tau)}^{\alpha, \tau} \leq f \ \& \ f_{\zeta_f(\tau)}^{\alpha, \tau} = f \text{ mod } J_\tau^\alpha\}.$$

(Note that if $\alpha = \delta$ then there is only one value of τ_ℓ, τ which we consider here: λ .) Let $T' \subseteq \bigcup_{i \leq \delta} \prod_{j < i} \lambda_j$ be a tree such that for $\gamma \leq \delta$:

$$T'_\gamma = \{f \in \prod_{i < \gamma} \lambda_i : f \restriction \alpha \in T_\alpha^0 \text{ for each limit } \alpha \leq \gamma\}.$$

Let

$$A = \{\zeta < \lambda : (\exists f \in \prod_{i \leq \delta} \lambda_i)[f_\zeta^{\delta, \lambda} \leq f \ \& \ f_\zeta^{\delta, \lambda} = f \bmod J_\lambda^\alpha \ \& \ (\forall i \leq \delta)(f \restriction i \in T'_\alpha)]\}$$

and for each $\zeta \in A$ let f_ζ^* be a function witnessing it. Now let $T \subseteq \bigcup_{i \leq \delta} \prod_{j < i} \lambda_j$

be a tree such that $T_\delta = \{f_\zeta^* : \zeta \in A\}$.

By the definition, T is a tree, but maybe it does not have enough levels? Let χ be a large enough regular cardinal. Take an increasing continuous sequence $\langle N_i : i \leq \theta \rangle$ of elementary submodels of $(\mathcal{H}(\chi), \in, <^*)$ such that

$$\begin{aligned} |N_i| &= \Upsilon = \theta + |\text{pcf}\{\lambda_\alpha : \alpha < \delta\}| < \lambda_0, \\ \Upsilon + 1 &\subseteq N_i \in N_{i+1}, \\ \text{all relevant things are in } N_0. \end{aligned}$$

Now we define $f^* \in \prod_{\alpha < \delta} \lambda_\alpha$ by

$$f^*(\alpha) = \sup(\bigcup_{i < \theta} N_i \cap \lambda_\alpha).$$

Similarly as in [Sh 355], pp 63–65, one proves that $f^* \restriction \alpha \in T_\alpha^0$ for each limit $\alpha \leq \delta$. Hence for some ζ we have $f^* = f_\zeta^{\delta, \lambda} \bmod J_\lambda^\delta$ and thus $\zeta \in A$. Consequently A is unbounded in λ .

By induction on $\alpha \leq \delta$ we prove that

($\tilde{\odot}$) if $f_\zeta \in T_\alpha$ (for $\zeta < \theta$) are pairwise distinct

then there are $Z \in [\theta]^\theta$ and $j < \alpha$ such that

$$(\forall \zeta_0, \zeta_1 \in Z)(\zeta_0 \neq \zeta_1 \Rightarrow [f_{\zeta_0} \restriction j = f_{\zeta_1} \restriction j \ \& \ f_{\zeta_0}(j) \neq f_{\zeta_1}(j)]).$$

If α is a non-limit ordinal then this is trivial. So suppose that α is limit, $\alpha < \delta$. Then for some $\tau_{\zeta, \ell} \in \text{pcf}\{\lambda_i : i < \alpha\}$, $\xi_{\zeta, \ell} < \tau_{\zeta, \ell}$, $n_\zeta < \omega$ (for $\zeta < \theta$, $\ell < n_\zeta$) we have

$$f_\zeta = \max\{f_{\xi_{\zeta, \ell}}^{\alpha, \tau_{\zeta, \ell}} : \ell < n_\zeta\}.$$

As $\theta > |\text{pcf}\{\lambda_\beta : \beta < \alpha\}|$ we may assume that $n_\zeta = n^*$, $\tau_{\zeta, \ell} = \tau_\ell$ and for each $\ell < n^*$ the sequence $\langle \xi_{\zeta, \ell} : \zeta < \theta \rangle$ is either constant or strictly increasing.

Now, the second case has to occur for some ℓ and we may follow similarly to 5.14.1 and then apply the inductive hypothesis. We are left with the case $\alpha = \delta$. So let $f_\zeta = f_{\beta_\zeta}^*$ for $\zeta < \delta$ and we continue as before (with λ for τ_ℓ). This ends the proof of the claim (note that the arguments showing that all the T_α^0 are not empty prove actually that the tree T has enough branches to satisfy our additional requirements).

Now let T be a tree is in the claim above. Let $\bar{\eta} = \langle \eta_\alpha : \alpha < \lambda \rangle \subseteq \lim_\delta(T)$ be the enumeration of $\{f_\zeta^* : \zeta \in A\}$ such that $\bar{\eta}$ is $<_{J_{<\lambda}}$ -increasing cofinal in $\prod_{i<\delta} \lambda_i / J_{<\lambda}$. By the assumption **(d)** for each $\eta \in T$ we find a marked Boolean algebra $(\mathbb{B}_\eta, \bar{y}_\eta)$ such that for every $i < \delta$ the sequence $\langle (\mathbb{B}_\eta, \bar{y}_\eta) : \eta \in T_i \rangle$ witnesses that $(\theta, \lambda_i, |T_i|) \in \mathcal{K}_{\text{smk}}$. These parameters determine a (δ, μ, λ) -constructor $\mathcal{C}$, so we have the respective Boolean algebra $\mathbb{B}^{\text{red}}(\mathcal{C})$ (and its counterpart $\mathbb{B}^{\text{green}}(\mathcal{C})$). To show that they have the required properties we follow exactly the proof that $(\theta, \lambda, \chi) \in \mathcal{K}_{\text{smk}}$, so we will present this only.

First note that by 5.13 the algebra $\mathbb{B}^{\text{red}}(\mathcal{C})$ has the θ -Knaster property. Now, let $\langle A_\zeta : \zeta < \chi \rangle \subseteq [\lambda]^\lambda$ be such that

$$\zeta_1 < \zeta_2 < \chi \quad \Rightarrow \quad |A_{\zeta_1} \cap A_{\zeta_2}| < \lambda.$$

Let $\bar{x}_\zeta = \langle x_\xi^{\text{red}} : \xi \in A_\zeta \rangle$ and let $\mathbb{B}_\zeta$ be the subalgebra of $\mathbb{B}^{\text{red}}(\mathcal{C})$ generated by $\bar{x}_\zeta$. We want to show that the sequence $\langle (\mathbb{B}_\zeta, \bar{x}_\zeta) : \zeta < \chi \rangle$ witnesses $(\theta, \lambda, \chi) \in \mathcal{K}_{\text{smk}}$. For this suppose that $\zeta_0 < \dots < \zeta_{n-1} < \chi$, $n < \omega$ and $\beta_{\varepsilon, \ell} \in A_{\zeta, \ell}$ are increasing with ε (for $\varepsilon < \lambda$, $\ell < n$) and without loss of generality with no repetition. We may assume that

$$(\forall \ell < n)(\forall \varepsilon < \lambda)(\beta_{\varepsilon, \ell} \notin \bigcup_{m \neq \ell} A_{\zeta_m}).$$

Further we may assume that for some $i^* < \delta$ and pairwise distinct $\eta_\ell \in T_{i^*}$ (for $\ell < n$) we have

$$(\forall \varepsilon < \lambda)(\forall \ell < n)(\eta_{\beta_{\varepsilon, \ell}} \restriction i^* = \eta_\ell).$$

Now we take $i \in [i^*, \delta)$ such that

$$(\forall \gamma < \lambda_i)(\exists^\lambda \varepsilon < \lambda)(\forall \ell < n)(\eta_{\beta_{\varepsilon, \ell}}(i) > \gamma)$$

(remember that each $\langle \eta_{\beta_{\varepsilon, \ell}} : \varepsilon < \lambda \rangle$ is $<_{J_{<\lambda}}$ -cofinal). Since $|T_i| < \lambda_i$ we can find $\nu_0, \dots, \nu_{n-1} \in T_i$ such that $\eta_\ell \sqsubseteq \nu_\ell$ and

$$(\forall \gamma < \lambda_i)(\exists^\lambda \varepsilon < \lambda)(\forall \ell < n)(\eta_{\beta_{\varepsilon, \ell}} \restriction i = \nu_\ell \ \& \ \eta_{\beta_{\varepsilon, \ell}}(i) > \gamma).$$

Consequently we may choose a sequence $\langle \langle \gamma_{\xi, \ell} : \ell < n \rangle : \xi < \lambda_i \rangle \subseteq \lambda_i$ such that $\xi < \gamma_{\xi, \ell}$ and

$$(\forall \xi < \lambda_i)(\exists^\lambda \varepsilon < \lambda)(\forall \ell < n)(\eta_{\beta_{\varepsilon, \ell}} \restriction (i+1) = \nu_\ell \hat{\ } \langle \gamma_{\xi, \ell} \rangle).$$

Now we use the choice of $(\mathbb{B}_{\nu_\ell}, \bar{y}_{\nu_\ell})$ (witnessing $(\theta, \lambda_i, |T_i|) \in \mathcal{K}_{\text{smk}}$) and we find $\xi_1 < \xi_2 < \lambda_i$ such that

$$(\forall \ell < n)(\mathbb{B}_{\nu_\ell} \models y_{\gamma_{\xi_1, \ell}}^{\nu_\ell} \cap y_{\gamma_{\xi_2, \ell}}^{\nu_\ell} = \mathbf{0}),$$

which allows us to find $\varepsilon_1 < \varepsilon_2 < \lambda$ such that for each $\ell < n$ the intersection $x_{\beta_{\varepsilon_1, \ell}} \cap x_{\beta_{\varepsilon_2, \ell}}$ is $\mathbf{0}$. $\blacksquare$

Conclusion 6.5 *If $\langle \mu_i : i \leq \kappa \rangle$ is a strictly increasing continuous sequence of strong limit singular cardinals such that $\kappa < \mu_0$, $2^{\mu_i} = \mu_i^+$, $\kappa < \theta = \text{cf}(\theta) < \mu_0$ and*

$$\{\alpha < \mu_i^+ : \text{cf}(\alpha) = \theta\} \in I[\mu_i^+]$$

then $(\theta, \mu_\kappa^+) \in \mathcal{K}_{\text{wmk}}$ and we may construct the respective Boolean algebras $\mathbb{B}^{\text{red}}, \mathbb{B}^{\text{green}}$. $\blacksquare$

Proposition 6.6 *Suppose that we have Boolean algebras $\mathbb{B}^{\text{red}}, \mathbb{B}^{\text{green}}$ such that*

*$\mathbb{B}^{\text{red}}$ satisfies the θ -Knaster condition
for each $n < \omega$ the free product $(\mathbb{B}^{\text{green}})^n$ satisfies the λ -cc
the free product $\mathbb{B}^{\text{red}} * \mathbb{B}^{\text{green}}$ fails the λ -cc.*

Then $(\theta, \lambda, \chi) \in \mathcal{K}_{\text{smk}}$, where $\chi = \lambda^+$ (or even if χ is such that there is an almost disjoint family $\mathcal{A} \subseteq [\lambda]^\lambda$ of size χ).

PROOF We have $y_\alpha \in (\mathbb{B}^{\text{red}})^+$ and $z_\alpha \in (\mathbb{B}^{\text{green}})^+$ for $\alpha < \lambda$ such that if $\alpha < \beta < \lambda$ then

$$\text{either } \mathbb{B}^{\text{red}} \models y_\alpha \cap y_\beta = \mathbf{0} \quad \text{or} \quad \mathbb{B}^{\text{green}} \models z_\alpha \cap z_\beta = \mathbf{0}.$$

Let $A_\zeta \in [\lambda]^\lambda$ (for $\zeta < \chi$) be pairwise almost disjoint sets. We want to show that the sequence

$$\langle (\mathbb{B}^{\text{red}}, \bar{y} \restriction A_\zeta) : \zeta < \chi \rangle$$

is a witness for $(\theta, \lambda, \chi) \in \mathcal{K}_{\text{smk}}$. So we are given $\zeta_0 < \zeta_1 < \dots < \zeta_{n-1} < \chi$ and sequences $\langle \alpha_{\varepsilon, \ell} : \varepsilon < \lambda \rangle \subseteq A_{\zeta_\ell}$. Then, for some $\varepsilon^* < \lambda$ we have

$$\varepsilon^* \leq \varepsilon < \lambda \quad \Rightarrow \quad \alpha_{\varepsilon, \ell} \notin \bigcup_{m \neq \ell} A_{\zeta_m}.$$

We should find $\varepsilon_1 < \varepsilon_2$ such that for all $\ell < n$

$$\mathbb{B}^{\text{red}} \models y_{\alpha_{\varepsilon_1, \ell}} \cap y_{\alpha_{\varepsilon_2, \ell}} = \mathbf{0}.$$

For this it is enough to find $\varepsilon^* < \varepsilon_1 < \varepsilon_2$ such that for $\ell < n$

$$\mathbb{B}^{\text{green}} \models z_{\alpha_{\varepsilon_1, \ell}} \cap z_{\alpha_{\varepsilon_2, \ell}} \neq \mathbf{0}.$$

But this we easily get from the fact that the free product $(\mathbb{B}^{\text{green}})^n$ satisfies the λ -cc. ■

Comment 6.7 *The proofs that the algebra $\mathbb{B}^{\text{green}}$ satisfies the λ -cc (see 4.3, 6.4) give that actually for each $n < \omega$ the product $(\mathbb{B}^{\text{green}})^n$ satisfies λ -cc. So it is reasonable to add it (though not needed originally).*

Comment 6.8 *The “ $\bar{\eta}$ is (strong-) J -cofinal for $(T, \bar{\lambda}, \bar{I})$ ” has easy consequences for the existence of colourings.*

Remark 6.9 For μ strong limit singular we may sometimes get a cofinal sequence of length $\lambda \in (\mu, 2^\mu]$ without $2^\mu = \mu^+$. By [Sh 430], section 5, if:

- (a) I_i is a χ_i -complete, $|I_i| = \tau_i$, χ_i regular
- (b) $\chi_i \leq \tau_i \leq (\chi_i)^{+n^*}$, $n^* < \omega$
- (c) $\text{tcf}(\prod_{i < \delta} (\chi_i)^{+\ell} / J) = \lambda$ for each $\ell \leq n^*$

then

- (α) there is a cofinal sequence in $\prod_{i < \delta} (\mathcal{P}(\lambda_i) / I_i) / J$, because

- (β) it has the true cofinality.

So if for arbitrarily large χ , $2^\chi = \chi^+$, $2^{\chi^+} = \chi^{++}$ then we have the ideal we want and maybe the pcf condition holds. so combining this and 6.10 below we get that there may be an example of our kind not because of GCH reasons, but still requiring some cardinal arithmetic assumptions.

Proposition 6.10 *Suppose that $\langle \lambda_i : i < \delta \rangle$ is a strictly increasing sequence of regular cardinals, I_i is a $(\prod_{j < i} \lambda_j)^+$ -complete ideal on λ_i (so $\prod_{j < i} \lambda_j < \lambda_i$) and $(\mathbb{B}_i, \bar{y}_i, I_i)$ is a λ_i -well marked Boolean algebra (for $i < \delta$).*

1. Assume that $\prod_{i < \delta} (I_i, \subseteq) / J$ has true cofinality λ . Then there exists a $(\theta, \text{not } \lambda)$ -Knaster marked Boolean algebra.

2. Suppose in addition that $h : \delta \longrightarrow \omega$ is a function such that

$$(\forall n < \omega)(h^{-1}[\{n\}] \in J^+)$$

and $I_i^{[h(i)]}$ are the product ideals on $(\lambda_i)^n$ (for $i < \delta$):

$$I_i^{[h(i)]} \stackrel{\text{def}}{=} \{B \subseteq (\lambda_i)^n : \neg(\exists^{I_i} \gamma_0) \dots (\exists^{I_i} \gamma_{h(i)-1})(\langle \gamma_\ell : \ell < h(i) \rangle \in B)\}.$$

Assume that

$$\lambda = \text{tcf}(\prod_{i < \delta} (I_i^{[h(i)]}, \subseteq) / J)$$

and that the $(\mathbb{B}_i, \bar{y}_i, I_i)$ satisfy the following requirement:

$(\ast)_{h(i)}$ if $B \subseteq (\text{dom}(\bar{y}_i))^{h(i)}$ is such that

$$(\exists^{I_i} \gamma_0) \dots (\exists^{I_i} \gamma_{h(i)})(\langle \gamma_\ell : \ell \leq h(i) \rangle \in B)$$

then there are $\gamma'_\ell, \gamma''_\ell < \lambda_i$ (for $\ell \leq h(i)$) such that for each ℓ

$$\mathbb{B}_i \models y_{i, \gamma'_\ell} \cap y_{i, \gamma''_\ell} = \mathbf{0}.$$

Then we can conclude that $((2^{|\delta|})^+, \lambda, \lambda^+) \in \mathcal{K}_{\text{smk}}$ and we have a pair of algebras $(\mathbb{B}^{\text{red}}, \mathbb{B}^{\text{green}})$ as in main theorem 4.4.

PROOF The main point here is that with our assumptions in hands we may construct a sequence $\langle \eta_\alpha : \alpha < \lambda \rangle \subseteq \prod_{i < \delta} \lambda_i$ which is quite stronger J -cofinal: it satisfies the requirement of 2.2(6)(b) weakened to the demand that the set there is not in the dual filter J^c . Of course this is still enough to carry our proofs and we may use such a sequence to build the right examples. 1). Let $\langle \langle A_i^\alpha : i < \delta \rangle : \alpha < \lambda \rangle$ witness the true cofinality. By induction on $\alpha < \lambda$ choose $\gamma_\alpha < \lambda$ and $\eta_\alpha \in \prod_{i < \alpha} \lambda_i$ such that

$$\langle \{\eta_\beta(i) : i < \delta\} \in \prod_{i < \delta} I_i,$$

if $\beta < \alpha$ then $\gamma_\beta < \gamma_\alpha$ and $(\forall^J i)(\eta_\beta(i) \in A_i^{\gamma_\alpha})$ and $\eta_\alpha(i) \notin A_i^{\gamma_\alpha}$.

For $\alpha = 0$ or α limit, first choose $\gamma_\alpha = \sup\{\gamma_{\alpha_1} + 1 : \alpha_1 < \alpha\}$ and then choose $\eta_\alpha(i)$ by induction on i .

For $\alpha = \alpha_1 + 1$ first note that

$$\langle \{\eta_{\alpha_1}(i) : i < \delta\} \in \prod_{i < \delta} I_i.$$

Hence for some $\gamma_\alpha^0 < \lambda$ we have

$$(\forall^J i)(\eta_{\alpha_1}(i) \in A_i^{\gamma_\alpha}).$$

Let $\gamma_\alpha = \max\{\gamma_{\alpha_1}, \gamma_\alpha^0\}$. Now choose $\eta_\alpha(i)$ by induction on i .

As I_i is $|T_i|^+$ -complete, clearly $\langle \eta_\alpha : \alpha < \lambda \rangle$ is J -cofinal for $(T, J, \bar{I})$ and 3.7, 3.8 give the conclusion.

2). The construction of $\bar{\eta}$ is in a sense similar to the one in the proof of 2.8, but we use our cofinality assumptions. We have a cofinal sequence in $\prod_{i < \delta} (I_i^{[h(i)]}, \subseteq) / J$:

$$\langle \langle A_i^\alpha : i < \delta \rangle : \alpha < \lambda \rangle.$$

For each A_i^α we have “Skolem functions” $f_{i,\ell}^\alpha$ for $\ell < h(i)$ (like in the proofs of 4.3.1, 5.4).

We define η_α by induction on $\alpha < \lambda$. In the exclusion list we put all substitutions by $\eta_{\gamma_0} \upharpoonright i, \dots, \eta_{\gamma_{\ell-1}} \upharpoonright i$ for $\gamma_k < \alpha$ to $f_{i,\ell}^\alpha$: each time we obtain a set in the ideal I_i and a member $\bar{A}$ of $\prod_{i < \delta} I_i$ such that if $(\forall^J i)(\eta(i) \notin A_i)$, $\eta \in \prod_{i < \delta} \lambda_i$ then η satisfies the demand. Eventually we have $|\alpha|^{<\omega}$ such elements of $\prod_{i < \delta} I_i$. Let them be $\{\bar{B}^{\alpha,\xi} : \xi \leq |\alpha| + \aleph_0\}$. Then for some γ_α

$$(\forall \xi < |\alpha| + \aleph_0)(\forall^J i < \delta)(B_i^{\alpha,\xi} \subseteq A_i^{\gamma_\alpha})$$

and similarly

$$(\forall \beta < \alpha)(\forall^J i < \delta)(\eta_\beta(i) \in A_i^{\alpha_i}).$$

Choose $\eta_\alpha \in \prod_{i < \delta} (\lambda_i \setminus A_i^{\gamma_\alpha})$. ■

Remark 6.11 One of the main tools used in this section are (variants of) the following observation:

if $(\mathbb{B}, \bar{y})$ is a λ -marked Boolean algebra such that $\mathbb{B}$ is θ -Knaster and if $\varepsilon(\alpha, \ell) < \lambda$ (for $\alpha < \lambda$, $\ell < n$) are pairwise distinct then for some $\alpha < \beta < \lambda$, for each $\ell < n$ we have

$\mathbb{B} \models y_{\varepsilon(\alpha, \ell)} \cap y_{\varepsilon(\beta, \ell)} = \mathbf{0}$
then $(\theta, \lambda, \lambda^+) \in \mathcal{K}_{\text{smk}}$.

Concluding Remarks 6.12 If μ is a strong limit singular cardinal, $\text{cf}(\mu) \leq \theta = \text{cf}(\theta) < \mu$ then, by the methods of [Sh 576], one may get consistency of

if an algebra $\mathbb{B}$ satisfies the θ -ccc
then it satisfies the μ^+ -Knaster condition.

One may formulate the following question now:

Question (mostly solved) 6.13 *Suppose that $\mathbb{B}$ is a Boolean algebra satisfying the θ -cc and λ is a regular cardinal between μ^+ and $(2^\mu)^+$. Does $\mathbb{B}$ satisfy the λ -Knaster condition?*

There a reasonable amount of information on consistency of the negative answer in the next section, though 6.13 is not fully answered there. But a real problem is the following.

Problem 6.14 *Assume $\lambda = \mu^+$, $\text{cf}(\mu) = \theta$ and μ is a strong limit cardinal. Suppose that an algebra $\mathbb{B}_0$ satisfies the λ -cc and an algebra $\mathbb{B}_1$ satisfies the θ^+ -cc.*

*Does the free product $\mathbb{B}_0 * \mathbb{B}_1$ satisfy the λ -cc? (is this consistent? see 5.15).*

Problem 6.15 *Is it consistent that*

each Boolean algebra with the $\aleph_1$ -Knaster property has the λ -Knaster property for every regular (uncountable) cardinal λ ?

7 Some consistency results

We had seen that without inner models with large cardinals we have a complete picture, e.g.:

($\aleph$) if $\theta = \text{cf}(\theta) > \aleph_0$, $\mathbb{B}$ is a Boolean algebra satisfying the θ -cc and λ is a regular cardinal such that

$$(\forall \tau < \lambda)(\tau^{<\theta} < \theta)$$

then the algebra $\mathbb{B}$ satisfies the λ -Knaster condition.

($\beth$) if $\theta = \text{cf}(\theta) > \aleph_0$, $\theta < \mu = \mu^{<\mu} < \lambda = \text{cf}(\lambda) < \chi = \chi^\lambda$

then there is a μ^+ -cc μ -complete forcing notion $\mathbb{P}$ of size χ such that

$$\Vdash_{\mathbb{P}} \text{“the } \theta\text{-cc implies the } \lambda\text{-Knaster property”}.$$

Moreover

($\beth$)⁺ if $\mu = \mu^{<\theta} < \lambda = \text{cf}(\lambda) \leq 2^\mu$ then the θ -cc implies the λ -Knaster property.

- ($\mathfrak{I}$) if $\theta = \text{cf}(\theta) < \mu$, μ is a strong limit singular cardinal, $\text{cf}(\mu) = \theta$
 then the θ^+ -cc does not imply the μ^+ -Knaster property (and even we have the product example).

In ($\mathfrak{I}$), if we allow (2^θ) -cc we may get even better conclusion. In this section we want to show, under a large cardinals hypothesis, the consistency of failure.

Proposition 7.1 *Assume that κ is a supercompact cardinal, $\kappa < \lambda = \text{cf}(\lambda)$. Let $\mathbb{B}$ be a Boolean algebra which does not have the λ -Knaster property. Then*

$$(\exists \theta)(\aleph_0 < \theta = \text{cf}(\theta) < \kappa \ \& \ \mathbb{B} \text{ does not have the } \theta\text{-Knaster property}).$$

PROOF Since κ is supercompact, for every second order formula ψ :

if $M \models \psi$
 then for some $N \prec M$, $|N| < \kappa$, $N \models \psi$

(see Kanamori and Magidor, [KnMg78]). ■

Proposition 7.2 1. *If $\aleph_0 < \lambda_0 < \lambda_1$ are regular cardinals such that*

$$(*)_{\lambda_0, \lambda_1} \text{ for every } x \in \mathcal{H}(\lambda_1^+) \text{ there is } N \prec (\mathcal{H}(\lambda_1^+), \in) \text{ such that } x \in N \text{ and } N \cong (\mathcal{H}(\lambda_0^+), \in)$$

then if a Boolean algebra $\mathbb{B}$ has the λ_0 -Knaster property then it has λ_1 -Knaster property (and $\mathbb{B} \models \lambda_0\text{-cc}$ implies $\mathbb{B} \models \lambda_1\text{-cc}$).

2. *The condition $(*)_{\lambda_0, \lambda_1}$ above holds if for some κ_0, κ_1 , $\kappa_0 < \lambda_0$, $\kappa_1 < \lambda_1$ we have:*

$$(\oplus) \text{ there is an elementary embedding } j : \mathbf{V} \longrightarrow M \text{ with the critical point } \kappa_0 \text{ and such that } j(\kappa_0) = \kappa_1, j(\lambda_0) = \lambda_1 \text{ and } M^{\lambda_1} \subseteq M.$$

3. *If κ_0 is a 2-huge cardinal (or actually less) and e.g. $\lambda_0 = \kappa_0^{+\omega+1}$ then for some $\lambda_1 = \kappa_1^{+\omega+1}$ the condition $(\oplus)$ above holds (we can assume GCH).*

PROOF Just check. ■

Proposition 7.3 *Assume that*

$$\mathbf{V} \models \text{“ GCH + there is 2-huge cardinal } > \theta = \text{cf}(\theta) \text{”}$$

(can think of $\theta = \aleph_0$). Then there is a θ -complete forcing notion $\mathbb{P}$ such that in $\mathbf{V}^{\mathbb{P}}$:

(a) *GCH holds*

(b) *if a Boolean algebra $\mathbb{B}$ has the θ^+ -Knaster property then it has the $\theta^{+\theta+1}$ -Knaster property*

(note that if $\aleph_\theta > \theta$ then $\theta^{+\theta+1} = \aleph_{\theta+1}$).

PROOF Similar to [LMSH 198]. ■

Chasing arrows what we use is

Proposition 7.4 *If $\mathbf{V} \models \text{GCH}$ (for simplicity), $\theta = \text{cf}(\theta) = \text{cf}(\mu) < \mu$, a Boolean algebra $\mathbb{B}$ does not satisfy the μ^+ -Knaster condition and $\mathbb{Q} = \text{Levy}(\theta, \mu)$ then $\mathbf{V}^{\mathbb{Q}} \models \text{“}\mathbb{B} \text{ does not have the } \theta^+ \text{-Knaster property”}$.* ■

8 More on getting the Knaster property

Our aim here is to get a ZFC result (under reasonable cardinal arithmetic assumptions) which implies that our looking for $(\kappa, \text{not } \lambda)$ -Knaster marked Boolean algebras near strong limit singular is natural. Bellow we discuss the relevant background. The proof relays on pcf theory (but only by quoting a simply stated theorem) and seems to be a good example of the applicability of pcf.

Theorem 8.1 *Assume $\mu = \mu^{<\aleph_\omega}$.*

1. *If a Boolean algebra $\mathbb{B}$ of cardinality $\leq 2^\mu$ satisfies the $\aleph_1$ -cc then $\mathbb{B}$ is μ -linked (see below).*
2. *If $\mathbb{B}$ is a Boolean algebra satisfying the $\aleph_1$ -cc then $\mathbb{B}$ has the λ -Knaster property for every regular cardinal $\lambda \in (\mu, 2^\mu]$.*

Where

Definition 8.2 1. *A Boolean algebra $\mathbb{B}$ is μ -linked if $\mathbb{B} \setminus \{0\}$ is the union of $\leq \mu$ sets of pairwise compatible elements.*

2. *A Boolean algebra $\mathbb{B}$ is μ -centered if $\mathbb{B} \setminus \{0\}$ is the union of $\leq \mu$ filters.*

Of course we can replace the $\aleph_1$ -cc, $\aleph_\omega$ by the κ -cc, $\aleph_\omega(\kappa)$ (see more later). The proof is self contained except relayence on a theorem quoted from [Sh 460].

Let us review some background. By [Sh:92], 3.1, if $\mathbb{B}$ is a κ -cc Boolean algebra of cardinality μ^+ and $\mu = \mu^{<\kappa}$ then $\mathbb{B}$ is μ -centered. The proof did not work for $\mathbb{B}$ of cardinality μ^{++} even if $2^\mu \geq \mu^{++}$ by [Sh 126], point being we consider three elements. But if $\mu = \mu^{<\mu} < \lambda^{<\lambda}$, for some μ^+ -cc μ -complete forcing notion $\mathbb{P}$ of cardinality λ , in $\mathbf{V}^{\mathbb{P}}$:

if $\mathbb{B}$ is a μ -c.c. Boolean algebra of cardinality $< \lambda$ the $\mathbb{B}$ is μ -centered

(follows from an appropriate axiom). Juhasz, Hajnal and Szentmiklossy [HaJuSz] continue this restricting themselves to μ -linked. Then proof can be carried for μ^{++} , and they continue by induction. However as in not few cases, the problem was for λ^+ , when $\text{cf}(\lambda) = \aleph_0$ so they assume

$\otimes$ if $\lambda \in (\mu, 2^\mu)$, $\text{cf}(\lambda) = \aleph_0$ then $\lambda = \lambda^{\aleph_0}$ and $\square_\lambda$

(on the square see Jensen [Jn]). This implies that if we start with $\mathbf{V} = \mathbf{L}$ and force, then the assumption $(\otimes)$ holds, so it is a reasonable assumption. Also they prove the consistency of the failure of the conclusion when $\otimes$ fails relying on [HJSh 249] (on a set system + graph constructed there) and on colouring of graphs (section 2 of [HaJuSz]), possibly $2^{\aleph_0} = \aleph_1$, $2^{\aleph_1} = \aleph_{\omega+1}$, $|\mathbb{B}| = 2^{\aleph_1}$, $\mathbb{B}$ satisfies the $\aleph_1$ -cc but is not $\aleph_1$ -linked, only $\aleph_2$ -linked.

This gives the impression of essentially closing the issue, and so I would have certainly thought some years ago, but this is not the case, exemplifying the danger of looking at specific cases. In fact, as we shall note in the end, their consistency result is best possible under our knowledge of relevant forcing methods. They use [HJSh 249] to have “many very disjoint sets” (i.e. $\langle X_\alpha : \alpha \in S \rangle$, $S \subseteq \{\delta < \aleph_{\omega+1} : \text{cf}(\delta) = \aleph_1\}$, $X_\alpha \subseteq \alpha = \sup X_\alpha$, and $\alpha \neq \beta \Rightarrow X_\alpha \cap X_\beta$ finite).

On pcf see [Sh:g]. Now, [Sh 460] has half jokingly a strong claim of proving GCH under reasonable reinterpretation. In particular [Sh 460] says there cannot be many strongly almost disjoint quite large sets, so this blocks reasonable extensions of [HaJuSz]. Now the main theorem of [Sh 460] enables us to carry the induction on $\lambda \in (\mu, 2^\mu]$ as in [Sh:92], 3.1, [HaJuSz], 3.x.

Proposition 8.3 *Suppose that:*

- (a) $\lambda > \theta = \text{cf}(\theta) \geq \kappa = \text{cf}(\kappa) > \aleph_0$
- (b) *there are a club E of λ and a sequence $\bar{\mathcal{P}} = \langle \mathcal{P}_\alpha : \alpha \in E \rangle$ (with $\alpha \in E \Rightarrow |\alpha| \mid \alpha$) such that*
 - (i) $\mathcal{P}_\alpha \subseteq [\alpha]^{<\kappa}$, $|\mathcal{P}_\alpha| \leq |\alpha|$ and $\bar{\mathcal{P}}$ is increasing continuous,

- (ii) if $X \subseteq \lambda$ has order type θ , then for some increasing $\langle \gamma_\varepsilon : \varepsilon < \kappa \rangle$ we have $\gamma_\varepsilon \in X$ and for each $\varepsilon < \kappa$ for some $\zeta \in (\varepsilon, \kappa)$ and $\alpha \leq \min(E \setminus \gamma_\varepsilon)$ we have $\{\gamma_\zeta : \zeta < \varepsilon\} \in \mathcal{P}_\alpha$,

(c) $\mathbb{B}$ is a Boolean algebra satisfying the κ -cc, $|\mathbb{B}| = \lambda$.

Then we can find a Boolean algebra $\mathbb{B}'$ and a sequence $\langle \mathbb{B}'_\alpha : \alpha \in E \rangle$ of subalgebras of $\mathbb{B}'$ such that

(α) $\mathbb{B} \subseteq \mathbb{B}' \subseteq \mathbb{B}^{\text{com}}$ (the completion)

(β) $\mathbb{B}' = \bigcup_{\alpha \in E} \mathbb{B}'_\alpha$, $|\mathbb{B}'_\alpha| \leq |\alpha| \neq \aleph_0$, $\langle \mathbb{B}'_\alpha : \alpha \in E \rangle$ increasing continuous in α

(γ) if $\alpha \in E$, $x \in \mathbb{B}' \setminus \{\mathbf{0}\}$ then for some $Y \subseteq \mathbb{B}'_\alpha \setminus \{\mathbf{0}\}$, $|Y| < \theta$ we have

if $y \in Y$ then $y \cap x = \mathbf{0}_{\mathbb{B}'}$ and

if $z \in \mathbb{B}'_\alpha$ is such that $z \cap x = \mathbf{0}_{\mathbb{B}'}$ then $z \leq \sup Y' \in \mathbb{B}'_\alpha$ for some $Y' \in [Y]^{<\kappa}$

(δ) if either $(*)_1$ or $(*)_2$ (see below) holds then we can add

Y generates the ideal $\{z \in \mathbb{B}'_\alpha : z \cap x = \mathbf{0}_{\mathbb{B}'}\}$

where

$(*)_1$ $(\forall \varepsilon < \theta)(|\varepsilon|^{<\kappa} < \theta)$

$(*)_2$ in (b) we can add:

if $X \subseteq \alpha$, $|X| < |\alpha|$ then for some τ , $\tau^{<\kappa} < \theta$ and $h : X \rightarrow \tau$ we have: if $Y \subseteq X$, $h \upharpoonright Y$ is constant then $Y \in \mathcal{P}_\alpha$.

PROOF Let χ be a large enough regular cardinal. Let $\mathbb{B} = \{x_\varepsilon : \varepsilon < \lambda\}$, let $\mathbb{B}^{\text{com}}$ be the completion of $\mathbb{B}$. We choose by induction on $\alpha \in E$ an elementary submodel N_α of $(H(\chi), \in, <^*_\chi)$ of cardinality $|\alpha|$, increasing continuous in α , such that $\mathbb{B}$, $\langle x_\varepsilon : \varepsilon < \lambda \rangle$, $\mathbb{B}^{\text{com}}$, $\bar{\mathcal{P}}$, λ , θ , κ belong to N_0 and $\langle N_\zeta : \zeta \leq \varepsilon \rangle \in N_{\varepsilon+1}$.

Note: if $\alpha \in \text{nacc}(E)$ then $\alpha \in N_\alpha$ hence $\mathcal{P}_\alpha \subseteq N_\alpha$.

Let

$$\mathbb{B}'_\alpha \stackrel{\text{def}}{=} N_\alpha \cap \mathbb{B}^{\text{com}}, \quad \mathbb{B}' = \bigcup_{\alpha \in E} \mathbb{B}'_\alpha.$$

We define by induction on $\alpha \in E$ a one-to-one function g_α from $\mathbb{B}'_\alpha$ onto α such that

$$\beta \in \alpha \cap E \Rightarrow g_\beta \subseteq g_\alpha, \text{ and } g_\alpha \text{ is the } <^*_\chi \text{-first such } g,$$

so $g_\alpha \in N_{\min(E \setminus (\alpha+1))}$. Let $g = \bigcup_{\alpha \in E} g_\alpha$. Thus g is a one-to-one function from $\mathbb{B}'$ onto λ . In the conclusion clauses (α) , (β) should be clear and let us prove clause (γ) . So let $\alpha \in E$, $x \in \mathbb{B}' \setminus \{\mathbf{0}\}$. We define $J = \{z \in \mathbb{B}'_\alpha : \mathbb{B}' \models "z \cap x = \mathbf{0}"\}$. Then J is an ideal of $\mathbb{B}'_\alpha$. We now try to choose by induction on $\varepsilon < \theta$, elements $y_\varepsilon \in J$ such that

- (i) y_ε is a member of $J \setminus \{\mathbf{0}_\mathbb{B}\}$
- (ii) there is no $u \in [\varepsilon]^{<\kappa}$ such that $y_\alpha \leq \sup_{\zeta \in u} y_\zeta \in \mathbb{B}'_\alpha$ (sup - in the complete Boolean algebra $\mathbb{B}^{\text{com}}$)
- (iii) under (i) + (ii), $g(y_\varepsilon)$ ($< \lambda$) is minimal (hence under (i) + (ii), $\beta_\varepsilon \stackrel{\text{def}}{=} \min\{\beta \leq \alpha : y_\varepsilon \in \mathbb{B}'_\beta\}$ is minimal).

If we are stuck for some $\varepsilon < \theta$, then for every $y \in J$ the condition (ii) fails (note that (iii) does not change at this point) i.e. there is a respective set u . So suppose y_ε is defined for $\varepsilon < \theta$. Clearly

$$\zeta < \varepsilon \Rightarrow g(y_\zeta) < g(y_\varepsilon)$$

and hence $\zeta < \varepsilon < \theta \Rightarrow \beta_\zeta \leq \beta_\varepsilon$, and $\zeta < \varepsilon \Rightarrow y_\zeta \neq y_\varepsilon$. Now apply clause (b)(ii) of the assumption to the set $X = \{\gamma_\varepsilon : \varepsilon < \theta\}$ to get a contradiction. ■

Proposition 8.4 *Suppose that*

- (a) $\lambda > \theta = \text{cf}(\theta) \geq \kappa = \text{cf}(\kappa) > \aleph_0$, and $\mu = \mu^{<\theta} \leq \lambda \leq 2^\mu$ and $(\forall \lambda < \theta)[|\lambda|^{<\kappa} < \theta]$,
- (b) as in 8.3 and either $(*)_1$ or $(*)_2$ of clause (δ) of 8.3,
- (c) $\mathbb{B}$ is a κ -cc Boolean algebra of cardinality λ ,
- (d) every subalgebra $\mathbb{B}' \subseteq \mathbb{B}^{\text{com}}$ of cardinality $< \lambda$ is μ -linked (see definition 8.2(1)).

Then $\mathbb{B}$ is μ -linked.

PROOF Let $\langle \mathbb{B}'_\alpha : \alpha \in E \rangle$ was as in the conclusion of 8.3. Without loss of generality we may assume that the set of elements $\mathbb{B}'_\alpha$ is α . Let for $\alpha \in E$, $h_\alpha : \mathbb{B}'_\alpha \setminus \{\mathbf{0}\} \rightarrow \mu$ be such that:

$$h_\alpha(x_1) = h_\alpha(x_2) \Rightarrow x_1 \cap x_2 \neq \mathbf{0}_\mathbb{B}.$$

For each $x \in \mathbb{B}' \setminus \mathbb{B}'_{\min(E)}$ let $\alpha(x) = \max\{\alpha \in E : x \notin \mathbb{B}'_\alpha\}$ (well defined as $\mathbb{B}' = \bigcup_{\alpha \in E} \mathbb{B}'_\alpha$ and $\langle \mathbb{B}'_\alpha : \alpha \in E \rangle$ is increasing continuous), and let $Y_{x,\alpha} \subseteq \mathbb{B}'_\alpha$ be such that $|Y_{x,\alpha}| < \theta$ and

$$Y_x \subseteq J_x \stackrel{\text{def}}{=} \{y \in \mathbb{B}'_\alpha : y \cap x = \mathbf{0}_\mathbb{B}\} \quad \text{and}$$

Y_x is cofinal in J_x (Y_x exists by 8.3, see clause (δ)).

Let us define $u_x^0 = \{0, \alpha(x)\}$ and Y_x^0 the subalgebra of $\mathbb{B}'$ generated by $\{x\}$, and $u_x^{n+1} = u_x^n \cup \{\alpha(y) : y \in Y_x^n\}$ and Y_x^{n+1} be the subalgebra of $\mathbb{B}'$ generated by

$$Y_x^n \cup \bigcup \{Y_{x_1,\alpha} : x_1 \in Y_x^n \text{ and } \alpha \in u_x^n\}.$$

Finally let $Y_x^\omega = \bigcup_{n < \omega} Y_x^n$. As θ is regular, $|Y_x^n| < \theta$ and as in addition θ is uncountable, $|Y_x^\omega| < \theta$. Let $u_x = \{\alpha(y) : y \in Y_x^\omega\}$. We can find $A_\zeta \subseteq \mathbb{B}' \setminus \{\mathbf{0}\}$ for $\zeta < \mu$ such that $\mathbb{B}' \setminus \{\mathbf{0}\} = \bigcup_{\zeta < \mu} A_\zeta$ and

($\tilde{\otimes}$) if $x_1, x_2 \in A_\zeta$, then there are one-to-one functions $f : Y_{x_1}^\omega \xrightarrow{\text{onto}} Y_{x_2}^\omega$ and $g : u_{x_1} \xrightarrow{\text{onto}} u_{x_2}$ such that:

- (i) f, g preserve the order,
- (ii) $f(x_1) = x_2$ and if $y \in Y_{x_1}^\omega$ then $g(\alpha(y)) = \alpha(f(y))$,
- (iii) if $\alpha \in u_{x_1}, y \in \mathbb{B}'_\alpha \cap Y_{x_1}^\omega$ then $h_\alpha(x_1) = h_{g(\alpha)}(f(y))$
- (iv) f is an isomorphism (of Boolean algebras)
- (v) g is the identity on $u_{x_1} \cap u_{x_2}$
- (vii) f is the identity on $Y_{x_1}^\omega \cap Y_{x_2}^\omega$

(Why? By [EK] or use $\langle \eta_x : x \in \mathbb{B}' \rangle, \eta_x \in {}^\mu 2$ with no repetitions.)

So it is enough to prove:

$$x_1, x_2 \in A_\zeta \Rightarrow x_1 \cap x_2 \neq \mathbf{0}_\mathbb{B}.$$

Let D_1 be an ultrafilter of $Y_{x_1}^\omega$ to which x_1 belongs, $D_2 =: \{f(y) : y \in Y_{x_2}^\omega\}$ (an ultrafilter on $Y_{x_2}^\omega$ to which x_2 belongs). It suffices to prove that for each $\alpha \in E$, $D_1 \cap \mathbb{B}_\alpha, D_2 \cap \mathbb{B}_\alpha$ generate non trivial filters on $\mathbb{B}_\alpha$. We do it by induction on α (note if $\alpha \leq \beta$ this holds for α provided it holds for β). If $\alpha \in u_{x_1} \cap u_{x_2}$ use clause (iii) of ($\tilde{\otimes}$) – note that this includes the case when $\alpha = 0$. For $\alpha \in \text{acc}(E)$ it follows by the finiteness of the condition. In the remaining case $\beta = \sup(E \cap \alpha) < \alpha$ and if $Y_{x_1}^\omega \cap \mathbb{B}'_\alpha \subseteq \mathbb{B}'_\beta, Y_{x_2}^\omega \cap \mathbb{B}'_\alpha \subseteq \mathbb{B}'_\beta$ this is trivial. So by symmetry we may assume that $\alpha \in u_{x_1} \setminus u_{x_2}$ and use the definition of Y_y for $y \in B_\alpha \cap Y_{x_1}^\omega \setminus \mathbb{B}_\beta$. $\blacksquare$

Proposition 8.5 *Assume $\mu = \mu^{<\aleph_\omega(\kappa)}$. Then for every $\lambda \in (\mu, 2^\mu]$ of cardinality $> \mu$, for every large enough regular $\theta < \aleph_\omega(\kappa)$ clause (b) of 8.3 holds.*

PROOF By [Sh 460], for every $\tau \in [\mu, \lambda)$ for some $\theta_\tau < \aleph_\omega(\kappa)$, we have:

($\tilde{\Theta}$) there is $\mathcal{P} = \mathcal{P}_\tau \subseteq [\tau]^{<\aleph_\omega(\kappa)}$ closed under subsets such that $|\mathcal{P}| \leq \tau$ and every $X \in [\tau]^{<\aleph_\omega(\kappa)}$ is the union of $< \theta_\tau$ members of members of $\mathcal{P}_\tau$.

Now as $\text{cf}(\lambda) > \mu$ for some $n < \omega$, the set

$$\Theta = \{\tau : \mu < \tau < \lambda, \theta_\tau \leq \aleph_\omega(\kappa)\}$$

is an unbounded subset of $\text{Card} \cap (\mu, \lambda)$. Let $\theta \geq (\aleph_{n+1}(\kappa))$ be regular. Now choose E club of λ such that $\alpha \in \text{nacc}(E) \Rightarrow |\alpha| \in \Theta$ and choose $\mathcal{P}_\alpha \subseteq [\alpha]^{<\kappa}$ increasing continuous with $\alpha \in E$, such that for $\alpha \in \text{nacc}(E)$, for every $X \in [\alpha]^\theta$, for some $h : X \rightarrow \aleph_n(\kappa)$, if $Y \subseteq X$, $|Y| < \kappa$ and $h \upharpoonright Y$ constant then $Y \in \mathcal{P}_\alpha$.

Now suppose $X \subseteq \lambda$, $\text{otp}(X) = \theta$, so let $X = \{\gamma_\varepsilon : \varepsilon < \theta\}$, β_ε increasing with ε ; let $\beta_\varepsilon = \min\{\alpha \in E : \gamma_\varepsilon < \beta\}$, so $\zeta < \varepsilon \Rightarrow \beta_\zeta \leq \beta_\varepsilon$ and $\beta_\varepsilon \in \text{nacc}(E)$ and there is $h_\varepsilon : \{\zeta : \zeta < \varepsilon\} \rightarrow \aleph_n(\kappa)$ such that for every $j < \aleph_n(\kappa)$,

$$u \in [\varepsilon]^{<\kappa} \ \& \ (h \upharpoonright u \text{ constant}) \Rightarrow \{\gamma_\zeta : \zeta \in u\} \in \mathcal{P}_{\beta_\varepsilon}.$$

Applying the Erdős–Rado theorem (i.e. $\theta \rightarrow (\aleph_n(\kappa)^+)^2_{\aleph_n(\kappa)}$) we get the desired result (the proof is an overkill). $\blacksquare$

Main Conclusion 8.6 *Suppose that κ is a regular uncountable cardinal, $\mu = \mu^{\aleph_\omega(\kappa)}$ and $\mathbb{B}$ is a Boolean algebra satisfying the κ -cc.*

1. *If $|\mathbb{B}| \leq 2^\mu$ then $\mathbb{B}$ is μ -linked*
2. *If λ is regular $\in (\mu, 2^\mu]$ then $\mathbb{B}$ satisfies the λ -Knaster condition.*

PROOF 1) We prove this by induction on $\lambda = |\mathbb{B}|$. If $|\mathbb{B}| \leq \mu$ this is trivial and if $\text{cf}(|\mathbb{B}|) \leq \mu$ this follows easily by the induction hypothesis. In other cases by 8.5, for some $\theta^* < \aleph_\omega(\kappa)$ for every regular $\theta \in (\theta^*, \aleph_\omega(\kappa))$, clause (b) of 8.3 holds. Choose $\theta = (\theta^{\aleph_0})^{++}$, so for this θ both clause (b) of 8.3 and $(*)_1$ of clause (δ) of 8.3 hold. Thus by claim 8.4 we can prove the desired conclusion for $\lambda = |\mathbb{B}|$.

2) Follows from part 1). $\blacksquare$

Proposition 8.7 1. *In 8.6 we can replace the assumption $\mu = \mu^{\aleph_\omega(\kappa)}$ by $\mu = \mu^{<\tau}$ if*

- $\otimes$ for every $\lambda \in (\mu, 2^\mu)$ of cardinality $> \mu$ for some $\theta = \text{cf}(\theta) \geq \kappa$ we have: clause (b) of 8.3 and $(*)_2$ of clause (δ) of 8.3 hold.
2. If $\lambda^* \in (\mu, 2^\lambda)$ and we want to have the conclusion of 8.6(1) with $|\mathbb{B}| = \lambda^*$ and 8.6(2) for λ^* -Knaster only then it suffice to restrict ourselves in $\otimes$ to $\lambda \leq \lambda^*$. ■

Proposition 8.8 In 8.3, if $(\forall \varepsilon < \theta)[|\varepsilon|^{<\kappa} < \theta]$ then we can weaken clause (ii) of assumption (b) to

(ii)' if $X \subseteq \lambda$ has order type θ then for some $\langle \gamma_\varepsilon : \varepsilon < \kappa \rangle$ we have: $\gamma_\varepsilon \in X$ and

$$(\forall \varepsilon < \kappa)(\exists \gamma \in X)(\exists \alpha \leq \min(E \setminus \gamma))(\{\gamma_\zeta : \zeta < \varepsilon\} \in \mathcal{P}_\alpha).$$

PROOF Let $X = \{j_\varepsilon : \varepsilon < \theta\}$ strictly increasing with ε , and let $\beta_\varepsilon = \min(E \setminus (j_\varepsilon + 1))$, so $\zeta < \varepsilon \Rightarrow \beta_\zeta \leq \beta_\varepsilon$. Let

$$\begin{aligned} e &\stackrel{\text{def}}{=} \{\varepsilon < \theta : \quad \varepsilon \text{ is a limit ordinal and} \\ &\quad \text{if } \varepsilon_1 < \varepsilon \text{ and } u \in [\varepsilon_1]^{<\kappa} \text{ and } \{j_\xi : \xi \in u\} \in \bigcup_{\zeta < \theta} \mathcal{P}_{\beta_\zeta} \\ &\quad \text{then } \{j_\varepsilon : \varepsilon \in u\} \in \bigcup_{\zeta < \varepsilon} \mathcal{P}_{\beta_\zeta}\}. \end{aligned}$$

Now e is a club of θ as (θ is regular and) $(\forall \varepsilon < \theta)[|\varepsilon|^{<\kappa} < \theta]$. So we can apply clause (ii)' to $X' = \{j_\varepsilon : \varepsilon \in e\}$, and get a subset $\{\gamma_\varepsilon : \varepsilon < \kappa\}$ as there, it is as required in clause (ii). ■

Proposition 8.9 1. Assume $\lambda > \theta = \text{cf}(\theta) \geq \kappa = \text{cf}(\kappa) > \aleph_0$. Then a sufficient condition for clause (b) + $(\delta)(*)_1$ of claim 8.3 is

- ($\tilde{\otimes}$) (a) $\lambda > \theta = \text{cf}(\theta)$
 (b) for arbitrarily large $\alpha < \lambda$ for some regular $\tau < \theta$ and $\lambda' < \lambda$, for every $\mathfrak{a} \subseteq \text{Reg} \cap |\alpha| \setminus \theta$ for some $\langle \mathfrak{b}_\varepsilon : \varepsilon < \varepsilon^* < \tau \rangle$ we have $\mathfrak{a} = \bigcup_{\varepsilon < \varepsilon^*} \mathfrak{b}_\varepsilon$ and $[\mathfrak{b}_\varepsilon]^{<\kappa} \subseteq J_{\leq \lambda'}[\mathfrak{a}]$ for every $\varepsilon < \varepsilon^*$.
 (c) $(\forall \varepsilon < \theta)[|\varepsilon|^{<\kappa} < \theta]$ or for every $\lambda' \in [\mu, \lambda]$, $\square_{\{\delta < \lambda' : \text{cf}(\delta) = \theta\}}$

2. Assume $\mu > \theta \geq \kappa = \text{cf}(\kappa) > \aleph_0$, a sufficient condition for clause (b) of 8.3 to hold is:

for every $\lambda \in [\mu, 2^\mu]$ of cofinality $> \mu$ for some $\theta' \leq \theta$, $\otimes_1$ holds (with θ' instead θ).

PROOF 1) By [Sh 430], [Sh 420], 2.6 or [Sh 513].

2) Follows. ■

Remark 8.10 So it is still possible that (assuming CH for simplicity)

$\otimes$ if $\mu = \mu^{\aleph_1}$, $\mathbb{B}$ is c.c.c. Boolean algebra, $|\mathbb{B}| \leq 2^\mu$ then $\mathbb{B}$ is μ -linked.

On the required assumption see [Sh 410], Hyp. 6.1(x).

Note that the assumptions of the form $\lambda \in I[\lambda]$ if added save us a little on pcf hyp. (we mention it only in 8.x). But if we are interested in the $[\kappa - \text{c.c.} \Rightarrow \lambda\text{-Knaster}]$, it can be waived.

References

- [EK] Ryszard Engelking and Monika Karłowicz. Some theorems of set theory and their topological consequences. *Fundamenta Math.*, **57**:275–285, 1965.
- [HJSh 249] Andras Hajnal, Istvan Juhász, and Saharon Shelah. Splitting strongly almost disjoint families. *Transactions of the American Mathematical Society*, **295**:369–387, 1986.
- [HaJuSz] Andras Hajnal, Istvan Juhasz, and Z. Szentmiklossy. On the structure of CCC partial orders. *Algebra Universalis*, **to appear**.
- [Jn] Ronald B. Jensen. The fine structure of the constructible hierarchy. *Annals of Math. Logic*, **4**:229–308, 1972.
- [Ju] Istvan Juhász. Cardinal functions. II. In *Handbook of set-theoretic topology*, pages 63–109. North Holland Publ. Co, 1984.
- [KnMg78] Akihiro Kanamori and Menachem Magidor. The evolution of large cardinal axioms in set theory. In *Higher Set Theory*, volume 669 of *Lecture Notes in Mathematics*, pages 99–275. Springer – Verlag, 1978.
- [LMSh 198] Jean Pierre Levinski, Menachem Magidor, and Saharon Shelah. Chang’s conjecture for $\aleph_\omega$. *Israel Journal of Mathematics*, **69**:161–172, 1990.
- [MgSh 204] Menachem Magidor and Saharon Shelah. When does almost free imply free? (For groups, transversal etc.). *Journal of the American Mathematical Society*, **7**(4), 1994.

- [M1] Donald Monk. *Cardinal Invariants of Boolean Algebras*. Lectures in Mathematics. ETH Zurich, Birkhauser Verlag, Basel Boston Berlin, 1990.
- [M2] Donald Monk. *Cardinal Invariants of Boolean Algebras*, volume 142 of *Progress in Mathematics*. Birkhäuser Verlag, Basel–Boston–Berlin, 1996.
- [Sh:g] Saharon Shelah. *Cardinal Arithmetic*, volume 29 of *Oxford Logic Guides*. Oxford University Press, 1994.
- [Sh:92] Saharon Shelah. Remarks on Boolean algebras. *Algebra Universalis*, **11**:77–89, 1980.
- [Sh:93] Saharon Shelah. Simple unstable theories. *Annals of Mathematical Logic*, **19**:177–203, 1980.
- [Sh 126] Saharon Shelah. On saturation for a predicate. *Notre Dame Journal of Formal Logic*, **22**:239–248, 1981.
- [Sh 345] Saharon Shelah. Products of regular cardinals and cardinal invariants of products of Boolean algebras. *Israel Journal of Mathematics*, **70**:129–187, 1990.
- [Sh 345a] Saharon Shelah. Basic: Cofinalities of small reduced products. In *Cardinal Arithmetic*, volume 29 of *Oxford Logic Guides*, chapter I. Oxford University Press, 1994.
- [Sh 355] Saharon Shelah. $\aleph_{\omega+1}$ has a Jonsson Algebra. In *Cardinal Arithmetic*, volume 29 of *Oxford Logic Guides*, chapter II. Oxford University Press, 1994.
- [Sh 410] Saharon Shelah. More on Cardinal Arithmetic. *Archive for Mathematical Logic*, **32**:399–428, 1993.
- [Sh 420] Saharon Shelah. Advances in Cardinal Arithmetic. In *Finite and Infinite Combinatorics in Sets and Logic*, pages 355–383. Kluwer Academic Publishers, 1993. N.W. Sauer et al (eds.).
- [Sh 430] Saharon Shelah. Further cardinal arithmetic. *Israel Journal of Mathematics*, **95**:61–114, 1996.
- [Sh 460] Saharon Shelah. The Generalized Continuum Hypothesis revisited. *Israel Journal of Mathematics*, **accepted**.

- [Sh 481] Saharon Shelah. Was Sierpiński right? III Can continuum-c.c. times c.c.c. be continuum-c.c.? *Annals of Pure and Applied Logic*, **78**:259–269, 1996.
- [Sh 513] Saharon Shelah. PCF and infinite free subsets. *Archive for Mathematical Logic*, **accepted**.
- [Sh 572] Saharon Shelah. Colouring and non-productivity of $\aleph_2$ -cc. *Annals of Pure and Applied Logic*, **84**:153–174, 1997.
- [Sh 576] Saharon Shelah. Categoricity of an abstract elementary class in two successive cardinals. *Israel Journal of Mathematics*, **submitted**.
- [So74] Robert M. Solovay. Strongly compact cardinals and the GCH. In *Proceedings of the Tarski Symposium, Berkeley 1971*, volume 25 of *Proceedings of Symposia in Pure Mathematics*, pages 365–372. A.M.S., 1974.